

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 December 2000 (14.12.2000)

PCT

(10) International Publication Number
WO 00/75114 A1

(51) International Patent Classification⁷: **C07D 215/26**,
A61K 31/395, A61P 43/00, C07D 405/12, C07C 233/43,
C07D 215/22, C07C 215/80, 311/08

David [GB/GB]; 13 Thelton Avenue, Broadbridge Heath,
Horsham, West Sussex RH12 3LS (GB).

(21) International Application Number: PCT/EP00/05058

(74) Agent: **BECKER, Konrad**; Novartis AG, Corporate
Intellectual Property, Patent & Trademark Department,
CH-4002 Basel (CH).

(22) International Filing Date: 2 June 2000 (02.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9913083.3 4 June 1999 (04.06.1999) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO,
NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR,
TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except AT, US*): **NO-
VARTIS AG** [CH/CH]; Schwarzwaldallee 215, CH-4058
Basel (CH).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for AT only*): **NOVARTIS-ERFINDUNGEN
VERWALTUNGSGESELLSCHAFT M.B.H.** [AT/AT];
Brunner Strasse 59, A-1230 Vienna (AT).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): **CUENOUD,
Bernard** [CH/GB]; Lodge 2, Novartis Horsham Research
Centre, Wimblehurst Road, Horsham, West Sussex RH12
5AB (GB). **BRUCE, Ian** [GB/GB]; 9 Bourns Court, Ayshe
Court Drive, Horsham, West Sussex RH13 5RL (GB).
FAIRHURST, Robin, Alec [GB/GB]; 5 Greenacres,
Ashington, West Sussex RH20 3LR (GB). **BEATTIE,**

Published:


- With international search report.
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: BETA2-ADRENOCEPTOR AGONISTS



(I)




(II)

(57) Abstract: Compounds of formula (I) in free or salt or solvate form, where Ar is a group of formula (II) Y is carbon or nitrogen and R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, X, n, p, q and r are as defined in the specification, their preparation and their use as pharmaceuticals, particularly for the treatment of obstructive or inflammatory airways diseases. The compounds of formula (I) in free, salt or solvate form, have β 2-adrenoreceptor agonist activity.

BETA2-ADRENOCEPTOR AGONISTS


This invention relates to organic compounds, their preparation and their use as pharmaceuticals.

The invention provides in one aspect a compound of formula



in free or salt or solvate form, where

Ar is a group of formula



R^1 is hydrogen, hydroxy, or alkoxy,

R^2 and R^3 are each independently hydrogen or alkyl,

R^4 , R^5 , R^6 and R^7 are each independently hydrogen, halogen, cyano, hydroxy, alkoxy, aryl, alkyl, alkyl substituted by one or more halogen atoms or one or more hydroxy or alkoxy groups, alkyl interrupted by one or more hetero atoms, alkenyl, trialkylsilyl, carboxy, alkoxycarbonyl, or $-\text{CONR}^{11}\text{R}^{12}$ where R^{11} and R^{12} are each independently hydrogen or alkyl, or R^4 and R^5 , R^5 and R^6 , or R^6 and R^7 together with the carbon atoms to which they are attached denote a carbocyclic or heterocyclic ring,

R^8 is halogen, $-\text{OR}^{13}$, $-\text{CH}_2\text{OR}^{13}$ or $-\text{NHR}^{13}$ where R^{13} is hydrogen, alkyl, alkyl interrupted by one or more hetero atoms, $-\text{COR}^{14}$, where R^{14} is hydrogen, $-\text{N}(\text{R}^{15})\text{R}^{16}$, alkyl or alkyl interrupted by one or more hetero atoms, or aryl and R^{15} and R^{16} are each independently hydrogen, alkyl or alkyl interrupted by one or more hetero atoms, or R^{13} is $-\text{C}(=\text{NH})\text{R}^{17}$,

-SOR¹⁷ or -SO₂R¹⁷ where R¹⁷ is alkyl or alkyl interrupted by one or more hetero atoms, and R⁹ is hydrogen, or R⁸ is -NHR¹⁸ where -NHR¹⁸ and R⁹, together with the carbon atoms to which they are attached, denote a 5- or 6- membered heterocycle,
R¹⁰ is -OR¹⁹ or -NHR¹⁹ where R¹⁹ is hydrogen, alkyl, alkyl interrupted by one or more hetero atoms, or -COR²⁰, where R²⁰ is -N(R²¹)R²², alkyl or alkyl interrupted by one or more hetero atoms, or aryl, and R²¹ and R²² are each independently hydrogen, alkyl or alkyl interrupted by one or more hetero atoms,
X is halogen or halomethyl or alkyl,
Y is carbon or nitrogen,
n is 1 or 2,
p is zero when Y is nitrogen or 1 when Y is carbon,
q and r are each zero or 1, the sum of q+r is 1 or 2; and
the carbon atom marked with an asterisk* has the R or S configuration, or a mixture thereof, when R¹ is hydroxy or alkoxy.

Terms used in this specification have the following meanings:

“Alkyl” denotes straight chain or branched alkyl, which may be, for example, C₁ to C₁₀ alkyl such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, straight or branched pentyl, straight or branched hexyl, straight or branched heptyl, straight or branched nonyl or straight or branched decyl. Preferably alkyl is C₁ to C₄ alkyl. Alkyl substituted by one or more halogen atoms or one or more hydroxy or alkoxy groups may be any of the above C₁ to C₁₀ alkyl groups substituted by one or more halogen, preferably fluorine or chlorine, atoms, by one or more hydroxy groups or by one or more C₁ to C₁₀, preferably C₁ to C₄, alkoxy groups.

“Alkyl interrupted by one or more hetero atoms” denotes straight chain or branched alkyl e.g. C₂ to C₁₀ alkyl, in which one or more pairs of carbon atoms are linked by -O-, -NR-, -S-, -S(=O)- or -SO₂-, where R is hydrogen or C₁ to C₁₀ (preferably C₁ to C₄) alkyl. Preferred such groups are alkoxyalkyl groups, preferably C₁-C₄-alkoxy-C₁-C₄-alkyl groups.

“Alkoxy” denotes straight chain or branched alkoxy and may be, for example, C₁ to C₁₀ alkoxy such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, or straight or branched pentoxy, hexyloxy, heptyloxy, octyloxy, nonyloxy or decyloxy. Preferably alkoxy is C₁ to C₄ alkoxy.

“Alkenyl” means straight chain or branched alkenyl, which may be unsubstituted or substituted, for example by one or more halogen atoms or one or more alkoxy groups, and which may be, for example, C₂ to C₁₀ alkenyl such as vinyl, 1-propenyl, 2-propenyl, 1-butenyl, isobutenyl, or straight or branched pentenyl, hexenyl, heptenyl, octenyl, nonenyl or decenyl. Preferred alkenyl is C₂ to C₄ alkenyl.

“Aryl” denotes unsubstituted or substituted aryl, e.g. unsubstituted phenyl or naphthyl, or phenyl or naphthyl substituted by one or more, e.g. 1 to 4, substituents selected from C₁-C₄-alkyl, hydroxy, C₁-C₄-alkoxy, halogen, or halo-C₁-C₄-alkyl. Preferably, aryl is unsubstituted phenyl or phenyl substituted by 1 or 2 substituents selected from C₁-C₄-alkyl or halogen.

“Alkylene” denotes straight chain or branched alkylene which may be, for example, C₁-C₁₀-alkylene such as methylene, ethylene, 1,2-propylene, 1,3-propylene, butylene, pentylene, hexylene, heptylene, octylene, nonylene or decylene. Preferably alkylene is C₁-C₄-alkylene.

“Alkenylene” denotes straight chain or branched alkenylene which may be, for example, C₂-C₁₀-alkenylene such as vinylene, propenylene, butenylene, pentenylene, hexenylene, heptenylene, octenylene, nonenylene or decenylene. Preferably alkenylene is C₂-C₄-alkenylene.

In formula I, n is 1 or 2, i.e. there are 2 or 4 CH₂ groups in the ring fused to the indicated benzene ring, so that ring is either a 5-membered or 7-membered ring.

The group Ar in formula II in which R⁸ is -NHR¹⁸ and -NHR¹⁸ and R⁹ together denote a 5- or 6- membered heterocycle may be, for example, a group in which Y is carbon, R⁸ is -NHR¹⁸ and -NHR¹⁸ and R⁹ together denote

a group of formula -NH-CO-R²³- where R²³ is an alkylene, alkenylene or alkyleneoxy group,

a group of formula -NH-SO₂-R²⁴ where R²⁴ is an alkyleneoxy group,

a group of formula -NH-R²⁵(COOR²⁶)- where R²⁵ is an alkylene or alkenylene group and R²⁶ is alkyl, or

a group of formula -NH-CO-NH- or -NH-CO-S-,

R¹⁰ is -OR¹⁹ where R¹⁹ is as hereinbefore defined,

X is alkyl,



p is 1, q is 1 and r is zero or 1.

The alkylene, alkenylene and alkyleneoxy groups preferably have 1 to 4 carbon atoms.



Preferred groups Ar of formula II in which R⁸ is -NHR¹⁸, and -NHR¹⁸ and R⁹ together denote a 5- or 6- membered heterocycle, include groups in which


Y is carbon, R⁸ is -NHR¹⁸ and -NHR¹⁸ and R⁹ together denote a group of formula -NH-CO-C(R²⁷)=C(R²⁸)- or -NH-CO-CH₂-O- or -NH-CO-CH₂- or -NH-SO₂-CH₂-O- or -NH-C(COOR²⁶)=CH- or -NH-CO-NH- or -NH-CO-S- where R²⁷ and R²⁸ are each independently hydrogen or C₁-C₄-alkyl and R²⁶ is C₁-C₄-alkyl, R¹⁰ is -OH, X is C₁-C₄-alkyl, p is 1, q is 1 and r is zero or 1.

More preferred groups Ar of formula II where R⁸ is -NHR¹⁸, and -NHR¹⁸ and R⁹ together denote a 5- or 6- membered heterocycle include those of the formulae




in which R²⁹, R³⁰ and R³¹ are each independently hydrogen or C₁-C₄-alkyl,






VI




VII

in which Z is -O-, -NH- or -S-.

The group Ar of formula II in which R^8 is halogen and R^9 is hydrogen may be, for example, a group of formula II in which Y is carbon, R^8 is halogen, preferably chlorine, R^9 is hydrogen, R^{10} is $-NHR^{18}$ where R^{18} is hydrogen or C_1 - C_4 -alkyl, preferably hydrogen or methyl, X is halogen or halomethyl, preferably chlorine or trifluoromethyl, and p, q and r are each 1. Preferred groups Ar among such groups include those of formulae





VIII




IX

The group Ar of formula II in which R^8 is $-OR^{13}$ and R^9 is hydrogen may be, for example, a group of formula II in which Y is carbon, R^8 is $-OR^{13}$ where R^{13} is hydrogen, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, $-COR^{14}$ where R^{14} is C_1 - C_4 -alkyl, C_6 - C_{10} -aryl or $-N(R^{15})R^{16}$ where R^{15} and R^{16} are each independently hydrogen or C_1 - C_4 -alkyl, R^{10} is $-OR^{19}$ or $-NHR^{19}$ where R^{19} is hydrogen, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, or $-COR^{20}$ where R^{20} is $-N(R^{21})R^{22}$, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl or C_6 - C_{10} -aryl and R^{21} and R^{22} are each independently hydrogen or C_1 - C_4 -alkyl, p and q are each 1 and r is zero. Preferred groups Ar among such groups include those of formulae




The group Ar of formula II in which R^8 is $-CH_2OR^{13}$ may be, for example, a group of formula II in which Y is carbon, R^8 is $-CH_2OR^{13}$ where R^{13} is hydrogen, C_1 - C_4 -alkyl, or C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, R^9 is hydrogen, R^{10} is $-OR^{19}$ where R^{19} is hydrogen, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl or R^{10} is $-NHR^{19}$ where R^{19} is hydrogen, C_1 - C_4 -alkyl or $-COR^{20}$ where R^{20} is C_1 - C_4 -alkyl, C_6 - C_{10} -aryl or $-N(R^{21})R^{22}$ where R^{21} and R^{22} are each independently hydrogen or C_1 - C_4 -alkyl, p and q are each 1 and r is zero; or a group of formula in which Y is nitrogen, R^8 is $-CH_2OR^{13}$ where R^{13} is hydrogen, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, R^{10} is $-OR^{19}$ where R^{19} is hydrogen, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy- C_1 -


C₄-alkyl, p and r are zero and q is 1. Preferred groups Ar among such groups include those of formulae



XII




XIII



XIV

The group Ar of formula II in which R⁸ is -NHR¹³ may be, for example, a group of formula II in which Y is carbon, R⁸ is -NHR¹³ where R¹³ is hydrogen, C₁-C₁₀ alkyl, C₁-C₁₀ alkyl interrupted by 1 to 3 hetero atoms, -COR¹⁴ where R¹⁴ is hydrogen, C₁-C₁₀-alkyl or C₁-C₁₀-alkyl interrupted by 1 to 3 hetero atoms, or R¹³ is -C(=NH)R¹⁷, -SOR¹⁷ or -SO₂R¹⁷ where R¹⁷ is C₁-C₁₀-alkyl or C₁-C₁₀-alkyl interrupted by 1 to 3 hetero atoms, R⁹ is hydrogen, R¹⁰ is -OR¹⁸ where R¹⁸ is hydrogen, C₁-C₄-alkyl or C₁-C₄-alkoxy-C₁-C₄ alkyl, p and q are each 1 and r is zero. Preferred groups Ar among such groups include those of formula



XV

especially those where R^{13} is hydrogen, C_1 - C_4 -alkyl, $-\text{COR}^{14}$ where R^{14} is hydrogen or C_1 - C_4 -alkyl, or R^{13} is $-\text{SO}_2\text{R}^{17}$ where R^{17} is C_1 - C_4 -alkyl.

Especially preferred groups Ar are those of formulae III, IV, V, XII and XV as hereinbefore defined.

The group R^1 in formula I may be, for example, hydrogen, hydroxy or C_1 - C_4 -alkoxy such as methoxy, ethoxy, isopropoxy, n-butoxy or tert-butoxy. Preferably, R^1 is hydroxy.

When R^1 is hydroxy or alkoxy, the carbon atom in formula I marked with an asterisk * preferably has the R configuration.

The groups R^2 and R^3 in formula I may be, for example, each independently hydrogen or C_1 - C_4 -alkyl, e.g. methyl or ethyl. In most of the preferred embodiments of the invention, R^2 is hydrogen and R^3 is hydrogen or methyl.

The groups R^4 , R^5 , R^6 and R^7 in formula I may be, for example, each independently hydrogen, chlorine, fluorine, chloromethyl, trifluoromethyl, hydroxy, C_1 - C_{10} -alkoxy, C_1 - C_{10} -alkyl, C_1 - C_{10} -alkyl interrupted by one or more oxygen or sulfur atoms or one or more NH, SO or SO_2 groups, C_2 - C_4 -alkenyl, trimethylsilyl, triethylsilyl, phenyl, carboxy, C_1 - C_4 -alkoxycarbonyl, $-\text{CONR}^{11}\text{R}^{12}$ (where R^{11} and R^{12} are each independently hydrogen or C_1 - C_4 -alkyl), or R^4 and R^5 , R^5 and R^6 or R^6 and R^7 , together with the carbon atoms to which they are attached, may denote a 5- or 6- membered carbocyclic ring, which is preferably a cycloaliphatic ring which is preferably saturated, or a 5- or 6- membered O- heterocyclic ring containing one or two oxygen atoms. Preferably, R^4 , R^5 , R^6 and R^7 are each hydrogen or are such that the benzene ring to which they are attached is symmetrically substituted, i.e. either (a) R^4 and R^7 are identical and R^5 and R^6 are identical or together denote a symmetrical ring, or (b) R^4 and R^5 together and R^6 and R^7 together denote identical rings. More preferably, R^4 and R^7 are identical and are each hydrogen, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy, and either R^5 and R^6 are identical and are each hydrogen, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy or C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, or R^5 and R^6 together denote $-(\text{CH}_2)_s-$ or $-\text{O}(\text{CH}_2)_t\text{O}-$ where s is 3 or 4 and t is 1 or 2.

Especially preferred compounds of the invention include compounds of formula I in which Ar is a group of formula III, IV, V, XII or XV, R^1 is hydroxy, R^2 and R^3 are hydrogen, and R^4 and R^7 are identical and are each hydrogen, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy, and either R^5

and R^6 are identical and are each hydrogen, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy or C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, or R^5 and R^6 together denote $-(CH_2)_4-$ or $-O(CH_2)_2O-$, in free or salt or solvate form. In such compounds, the carbon atom in formula I marked with an asterisk * preferably has the R configuration. Specific especially preferred compounds are those described in the Examples hereinafter.


The compounds of formula (I) are capable of forming acid addition salts, particularly pharmaceutically acceptable acid addition salts. Pharmaceutically acceptable acid addition salts of the compound of formula I include those of inorganic acids, for example, hydrohalic acids such as hydrofluoric acid, hydrochloric acid, hydrobromic acid or hydroiodic acid, nitric acid, sulfuric acid, phosphoric acid; and organic acids such as formic acid, acetic acid, propionic acid, butyric acid, benzoic acid, o-hydroxybenzoic acid, p-hydroxybenzoic acid, p-chlorobenzoic acid, diphenylacetic acid, triphenylacetic acid, 1-hydroxynaphthalene-2-carboxylic acid, 3-hydroxynaphthalene-2-carboxylic acid, aliphatic hydroxy acids such as lactic acid, citric acid, tartaric acid or malic acid, dicarboxylic acids such as fumaric acid, maleic acid or succinic acid, and sulfonic acids such as methanesulfonic acid or benzenesulfonic acid. These salts may be prepared from compounds of formula I by known salt-forming procedures.

Suitable solvates are pharmaceutically acceptable solvates, preferably hydrates.


The present invention also provides a process for the preparation of compounds of formula I in free or salt or solvate form. They can be prepared by a process comprising:

(a) for the preparation of a compound where R^1 is hydroxy, either

(i) reacting a compound of formula




with a compound of formula



where Ar^1 is Ar as hereinbefore defined or a protected form thereof, R^2 , R^3 , R^4 , R^5 , R^6 , R^7 and n are as hereinbefore defined and R^{32} is hydrogen, or an amine-protective group, or

(ii) reducing a compound of formula



where Ar^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as hereinbefore defined, to convert the indicated keto group into $-\text{CH}(\text{OH})-$; or

(b) for the preparation of a compound where R^1 is hydrogen, reducing a corresponding compound of formula I where R^1 is hydroxy; or

(c) for the preparation of a compound of formula I where R^1 is alkoxy, either (i) O-alkylating a corresponding compound of formula I where R^1 is hydroxy or (ii) reacting a corresponding compound having a leaving moiety instead of R^1 with an alcohol of formula R^1H where R^1 is alkoxy;

and, optionally, converting a resultant compound of formula I in protected form into a corresponding compound in unprotected form;

and recovering the resultant compound of formula I in free or salt or solvate form.

Process variant (a)(i) may be carried out using known procedures for epoxide-amine reactions. It is conveniently carried out without a solvent or in an inert solvent, for example

a hydrocarbon such as toluene or an alcohol such as n-butanol. The reaction temperature is conveniently from 25°C to 200°C, preferably from 80°C to 150°C. The temperature may be achieved by conventional heating or by microwave irradiation.


Process variant (a)(ii) may be carried out using conventional methods, for example by reaction with sodium borohydride under conventional conditions.

Process variant (b) may be carried out using known procedures for reduction of secondary alcohols to hydrocarbons. Process variant (c)(i) may be carried out using known procedures for O-alkylation, for example by reaction with an alkylating agent such as an alkyl halide under known conditions. Process variant (c)(ii) may be effected using known procedures for benzylic displacement reactions, the leaving moiety being e.g. tosylate, mesylate, halogen or hydroxy.

Compounds of formula I in free form may be converted into salt or solvate forms, and vice versa, in a conventional manner.


Compounds of the invention can be recovered from the reaction mixture and purified in a conventional manner. Isomers, such as enantiomers, may be obtained in a conventional manner, e.g. by fractional crystallization or asymmetric synthesis from corresponding asymmetrically substituted, e.g. optically active, starting materials.

Compounds of formula XVI are known compounds or can be prepared by processes analogous to those used for the preparation of the known compounds, for example the procedures described in Journal of Medicinal Chemistry 1987, 30, 1563-1566. Compounds of formula XVI in which the carbon atom indicated by the asterisk * is chiral may be prepared from a compound of formula



where Ar¹ and R² are as hereinbefore defined and L is a leaving atom or group, as described in WO95/25104.

Compounds of formula XVI may alternatively be prepared by epoxidation of a compound of formula




where Ar^1 and R^2 are as hereinbefore defined, using conventional procedures, such as those used in the Examples hereinafter.

Compounds of formula XX are known or may be prepared by methods analogous to those used for the preparation of known compounds, for example those used in the Examples hereinafter.


Compounds of formula XVII are known or may be prepared by methods analogous to those used for the preparation of the known compounds. R^{32} as an amine-protective group in formula XVII may be a known such group, for example as described in Protective Groups in Organic Synthesis, T.W.Greene, P.G.M. Wuts, John Wiley & Sons Inc, Second Edition, 1991, preferably benzyl or trifluoroacetyl.

For example, compounds of formula XVII, where R^3 is hydrogen, may be prepared by reducing an oxime of formula




where R^4 , R^5 , R^6 , R^7 and n are as hereinbefore defined. The reduction may be carried out by conventional methods for reducing oximes to amines. For example, the reduction may be carried out by catalytic hydrogenation, preferably using palladium on charcoal as the catalyst. The hydrogenation may be effected using known procedures, for example as described by R.D. Sindelar et al, J. Med. Chem. (1982), 25(7), 858-864. Oximes of formula XXI may be prepared as described by Sindelar et al, op.cit., or by analogous procedures.

Compounds of formula XVII where R^4 and R^7 are hydrogen can be prepared by reacting a compound of formula



with a compound of formula



where R^3 , R^5 , R^6 , R^{32} and n are as hereinbefore defined. The reaction may be carried out in the presence of a catalyst such as tris(triphenylphosphine)rhodium chloride. The reaction temperature may be, for example, from 60 to 120°C. The reaction is conveniently carried out in an inert solvent, for example ethanol, when the reaction temperature is conveniently about the reflux temperature of the solvent. The reaction may be carried out using known procedures, for example as described in WO96/23760. Where R^5 and R^6 are trialkylsilyl, the reaction between the compounds of formulae XXII and XXIII may be carried out in the presence of a metal carbonyl complex catalyst, for example using the procedure described by K.P.C. Vollhardt and R. Hillard, J. Am. Chem. Soc. 1977, 99(12), 4058, or an analogous procedure. Compounds of formula XXII may be prepared as described in WO96/23760 or by analogous procedures. Compounds of formula XXIII are known or may be prepared by known procedures.

Compounds of formula XVII where R^3 is alkyl, particularly methyl, and n is 1 may be prepared by amination of the corresponding 2-alkyl-indan-1-one using ammonia and K_3FeCN_6 , e.g. by the procedure of Fornum and Carlson, Synthesis 1972, 191.

Compounds of formula XVII as hereinbefore defined where R^4 , R^5 , R^6 and R^7 are such that the benzene ring to which they are attached is symmetrically substituted are novel, other than the compounds where R^4 , R^5 , R^6 , R^7 and R^{30} are each hydrogen, where R^4 and R^7 are methyl or methoxy when R^5 , R^6 and R^{30} are each hydrogen, and where R^4 , R^7 and R^{30} are hydrogen when R^5 and R^6 are each hydroxy, fluorine or chlorine. In particular, preferred intermediates of formula XVII are novel where (i) R^4 and R^7 are each hydrogen and R^5 and R^6 are either each C_2 - C_4 -alkyl, C_2 - C_4 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl or R^5 and R^6 together denote $-(\text{CH}_2)_s-$ or $-\text{O}(\text{CH}_2)_t\text{O}-$ where s is 1 to 4 and t is 1 or 2; or (ii) R^4 and R^7 are each C_2 - C_4 -alkyl or C_2 - C_4 -alkoxy and R^5 and R^6 are either each hydrogen, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy and R^5 and R^6 are either each hydrogen, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, or C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl or R^5 and R^6 together denote $-(\text{CH}_2)_s-$ or $-\text{O}(\text{CH}_2)_t\text{O}-$ where s is 1 to 4 and t is 1 or 2.

Compounds of formula XVIII are novel compounds which may be prepared by reaction of a compound of formula



where Ar¹ is as hereinbefore defined and Hal is a halogen atom, preferably chlorine or bromine, with a compound of formula XVII as hereinbefore defined. The reaction may be carried out using conventional procedures, for example those described by Yoshizaki et al, J. Med. Chem (1976), 19(9), 1138-42.

Where desired, the protection of any reactive group may be carried out at any appropriate stage in the above processes. The protecting group is suitably one used conventionally in the art and may be introduced and removed using conventional procedure. For example, when a hydroxy group in Ar¹ is protected by a benzyl group, the latter may be removed by catalytic hydrogenation in the presence of palladium on charcoal using conventional procedures, such as those used hereinafter in the Examples.

Compounds of formula I in free, salt or solvate form are useful as pharmaceuticals. Accordingly the invention also provides a compound of formula I in free, salt or solvate form for use as a pharmaceutical. The compounds of formula I in free, salt or solvate form, hereinafter referred to alternatively as "agents of the invention", have good β 2-adrenoreceptor agonist activity. The β 2 agonist activity, onset of action and duration of action of the agents of the invention may be tested using the guinea pig tracheal strip in vitro assay according to the procedure of R.A. Coleman and A.T. Nials, J.Pharmacol. Methods (1989), 21(1), 71-86. The binding potency and selectivity for the β 2-adrenoreceptor relative to the β 1-adrenoreceptor can be measured by a classical filtration binding assay according to the procedure of Current Protocols in Pharmacology (S.J.Enna(editor-in-chief) et al, John Wiley & Son, Inc, 1998), or by cAMP determination in cells expressing β 2- or β 1-adrenoceptor, according to the procedure of B. January et al, British J. Pharmacol. 123: 701-711 (1998).

The agents of the invention commonly have a rapid onset of action and have a prolonged stimulating action on the β 2-adrenoreceptor, compounds of the Examples hereinbelow having Ki (β 2) values of the order of 0.1 to 1000 nM, having durations of action of the

order of 1 to greater than 12 hours, and having binding selectivities for the β 2-adrenoreceptor relative to the β 1-adrenoreceptor from 1.5 to 500. For example, the compounds of Examples 1, 2, 4, 5, 6, 8, 27 and 29 have β 2 and β 1 binding potencies, measured by cAMP determination in cells expressing β 2- and β 1- adrenoreceptors, represented by EC_{50} values (β 2/ β 1) (in nM) of 0.92/9.52, 0.23/1.25, 6.07/14.5, 0.79/6.10, 0.3/3.60, 0.57/8.46 and 0.012/0.5 respectively. The compounds of Examples 2, 4, 5, 27 and 29 have T(50%) times (in minutes) of >400 at 71nM concentration, 82 at 100 nM, 444 at 100nM, 222 at 1.0nM and 279 at 10nM respectively in the guinea pig tracheal strip assay, where T(50%) is the time for inhibition of contraction to decay to 50% of its maximum value.

Having regard to their β 2 agonist activity, the agents of the invention are suitable for use in the treatment of any condition which is prevented or alleviated by activation of the β 2-adrenoreceptor. In view of their long acting selective β 2 agonist activity, the agents of the invention are useful in the relaxation of bronchial smooth muscle and the relief of bronchoconstriction. Relief of bronchoconstriction can be measured in models such as the in vivo plethysmography models of Chong et al, J. Pharmacol.Toxicol. Methods 1998, 39, 163-168, Hammelmann et al, Am. J. Respir. Crit. Care Med., 1997, 156, 766-775 and analogous models. The agents of the invention are therefore useful in the treatment of obstructive or inflammatory airways diseases. In view of their long duration of action, it is possible to administer the agents of the invention once-a-day in the treatment of such diseases. In another aspect, agents of the invention commonly exhibit characteristics indicating a low incidence of side effects commonly encountered with β 2 agonists such as tachycardia, tremor and restlessness, such agents accordingly being suitable for use in on demand (rescue) treatment as well as prophylactic treatment of obstructive or inflammatory airways diseases.

Treatment of a disease in accordance with the invention may be symptomatic or prophylactic treatment. Inflammatory or obstructive airways diseases to which the present invention is applicable include asthma of whatever type or genesis including both intrinsic (non-allergic) asthma and extrinsic (allergic) asthma. Treatment of asthma is also to be understood as embracing treatment of subjects, e.g. of less than 4 or 5 years of age, exhibiting wheezing symptoms and diagnosed or diagnosable as "wheezy infants", an established patient category of major medical concern and now often identified as incipient

or early-phase asthmatics. (For convenience this particular asthmatic condition is referred to as "wheezy-infant syndrome".)

Prophylactic efficacy in the treatment of asthma will be evidenced by reduced frequency or severity of symptomatic attack, e.g. of acute asthmatic or bronchoconstrictor attack, improvement in lung function or improved airways hyperreactivity. It may further be evidenced by reduced requirement for other, symptomatic therapy, i.e. therapy for or intended to restrict or abort symptomatic attack when it occurs, for example anti-inflammatory (e.g. corticosteroid) or bronchodilatory. Prophylactic benefit in asthma may in particular be apparent in subjects prone to "morning dipping". "Morning dipping" is a recognised asthmatic syndrome, common to a substantial percentage of asthmatics and characterised by asthma attack, e.g. between the hours of about 4 to 6 am, i.e. at a time normally substantially distant from any previously administered symptomatic asthma therapy.

Other inflammatory or obstructive airways diseases and conditions to which the present invention is applicable include adult respiratory distress syndrome (ARDS), chronic obstructive pulmonary or airways disease (COPD or COAD), including chronic bronchitis, or dyspnea associated therewith, emphysema, as well as exacerbation of airways hyperreactivity consequent to other drug therapy, in particular other inhaled drug therapy. The invention is also applicable to the treatment of bronchitis of whatever type or genesis including, e.g., acute, arachidic, catarrhal, croupus, chronic or phthinoid bronchitis. Further inflammatory or obstructive airways diseases to which the present invention is applicable include pneumoconiosis (an inflammatory, commonly occupational, disease of the lungs, frequently accompanied by airways obstruction, whether chronic or acute, and occasioned by repeated inhalation of dusts) of whatever type or genesis, including, for example, aluminosis, anthracosis, asbestosis, chalicosis, ptilosis, siderosis, silicosis, tabacosis and byssinosis.

Having regard to their β_2 agonist activity, the agents of the invention are also useful in the treatment of a condition requiring relaxation of smooth muscle of the uterus or vascular system. They are thus useful for the prevention or alleviation of premature labour pains in pregnancy. They are also useful in the treatment of chronic and acute urticaria, psoriasis, allergic conjunctivitis, actinitis, hay fever, and mastocytosis.

The agents of the invention are also useful as co-therapeutic agents for use in conjunction with anti-inflammatory or bronchodilatory drug substances, particularly in the treatment of obstructive or inflammatory airways diseases such as those mentioned hereinbefore, for example as potentiators of therapeutic activity of such drugs or as a means of reducing required dosaging or potential side effects of such drugs. An agent of the invention may be mixed with the anti-inflammatory or bronchodilatory drug in a fixed pharmaceutical composition or it may be administered separately, before, simultaneously with or after the anti-inflammatory or bronchodilatory drug. Such anti-inflammatory drugs include steroids, in particular glucocorticosteroids such as budesonide, beclamethasone, fluticasone or mometasone, and dopamine receptor agonists such as cabergoline, bromocriptine or ropinirole. Such bronchodilatory drugs include anticholinergic or antimuscarinic agents, in particular ipratropium bromide, oxitropium bromide and tiotropium bromide. Combinations of agents of the invention and steroids may be used, for example, in the treatment of COPD or, particularly, asthma. Combinations of agents of the invention and anticholinergic or antimuscarinic agents or dopamine receptor agonists may be used, for example, in the treatment of asthma or, particularly, COPD.

In accordance with the foregoing, the present invention also provides a method for the treatment of an obstructive or inflammatory airways disease which comprises administering to a subject, particularly a human subject, in need thereof a compound of formula I, or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore described. In another aspect, the invention provides a compound of formula I, or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore described for use in the preparation of a medicament for the treatment of an obstructive or inflammatory airways disease.

The agents of the invention may be administered by any appropriate route, e.g. orally, for example in the form of a tablet or capsule; parenterally, for example intravenously; topically to the skin, for example in the treatment of psoriasis; intranasally, for example in the treatment of hay fever; or, preferably, by inhalation, particularly in the treatment of obstructive or inflammatory airways diseases.

In a further aspect, the invention also provides a pharmaceutical composition comprising a compound of formula I in free form or in the form of a pharmaceutically acceptable salt or solvate thereof, optionally together with a pharmaceutically acceptable diluent or carrier therefor. Such compositions may be prepared using conventional diluents or excipients and techniques known in the galenic art. Thus oral dosage forms may include tablets and

capsules. Formulations for topical administration may take the form of creams, ointments, gels or transdermal delivery systems, e.g. patches. Compositions for inhalation may comprise aerosol or other atomizable formulations or dry powder formulations.

The invention also includes (A) a compound of formula I as hereinbefore described in free form, or a pharmaceutically acceptable salt or solvate thereof, in inhalable form; (B) an inhalable medicament comprising such a compound in inhalable form together with a pharmaceutically acceptable carrier in inhalable form; (C) a pharmaceutical product comprising such a compound in inhalable form in association with an inhalation device; and (D) an inhalation device containing such a compound in inhalable form.

Dosages employed in practising the invention will of course vary depending, for example, on the particular condition to be treated, the effect desired and the mode of administration. In general, suitable daily dosages for administration by inhalation are of the order of from 1 to 5000 µg.

The invention is illustrated by the following Examples. Compounds used in the Examples are prepared as follows:

Intermediate 1 - 5,6-Diethyl-inden-2-ylamine hydrochloride

Preparation 1 - 3-chloro-1-(3,4-diethylphenyl)- 1-propanone

1,2-Diethylbenzene (10.9 g, 74.6 mmol) and propionyl chloride (9.7 g, 74.6 mmol) are added dropwise to AlCl₃ (22.3 g, 167.8 mmol) in nitromethane (75 mL) over 30 min. The reaction mixture is stirred at room temperature for 2 hours, after which 70 g of ice and 14 mL concentrated sulphuric acid are added. The aqueous phase is extracted with ether, and the combined organic phases extracted with 2N HCl and saturated aqueous NaCl. The organic phase is further treated with activated charcoal, magnesium sulphate, and filtered, and the solvent removed *in vacuo*.

¹H-NMR (CDCl₃) ppm: 7.8 (1H, s, Ar); 7.7 (1H, d, Ar); 7.2 (1H, d, Ar); 3.9 (2H, t, CH₂); 3.4 (2H, t, CH₂); 2.8 (4H, q, CH₂CH₃); 1.2 (6H, m, CH₃).

Preparation 2 - 2,3-dihydro-5,6-diethyl-1H-inden-1-one

3-chloro-1-(3,4-diethylphenyl)- 1-propanone (15.5 g) is dissolved in 66 mL concentrated sulphuric acid and heated to 90 °C for 4 hours. The reaction mixture is cooled, ice (70 g) is added, and the aqueous solution extracted twice with toluene. The organic layer is washed

with sodium bicarbonate, saturated aqueous NaCl, and treated with activated charcoal and magnesium sulphate. After filtration, the solvent is removed *in vacuo*. The product is purified by flash column chromatography (silica, hexane / ethylacetate 10:1), and further crystallised in hexane.

¹H-NMR (CDCl₃) ppm: 7.6 (1H, s, Ar); 7.3 (1H, d, Ar); 3.1 (2H, m, CH₂); 2.7 (6H, m, CH₂+CH₂CH₃); 1.2 (6H, m, CH₃).

Preparation 3 - 5,6-Diethyl- 3-oxime-1H-indene-1,2(3H)-dione

2,3-Dihydro-5,6-diethyl-1H-inden-1-one (5 g, 26 mmol) in methanol (75 mL) is brought to 40 °C, n-butyl nitrite (3.0 g, 28.6 mmol) is added dropwise, followed by the addition of concentrated HCl (1.25 mL). After 1 hour, the reaction is brought to room temperature and the precipitated product filtered off, washed with ice-cold methanol and dried.

¹H-NMR (d₆-DMSO) ppm: 12.6 (1H, s, OH); 7.4 (1H, s, Ar); 7.3 (1H, d, Ar); 3.6 (2H, s, CH₂); 2.6 (4H, m, CH₂CH₃); 1.1 (6H, m, CH₃).

Preparation 4 - 5,6-Diethyl-indan-2-ylamine hydrochloride

5,6-Diethyl- 3-oxime-1H-indene-1,2(3H)-dione (4.5 g) is added to a mixture of acetic acid (150 mL), and concentrated sulphuric acid (4.5 mL). Pd/C 5% (1.5 g) is added, the reaction mixture degassed with nitrogen, and hydrogenated for 5 hours. The catalyst is then removed by filtration, the pH brought to pH 10 with 4M NaOH, and the solution extracted with chloroform. The organic phase is dried with magnesium sulphate, and the solvent removed *in vacuo*. The residue is redissolved in a minimum amount of ether, and HCl saturated ether added. The white precipitate is filtered and dried to yield the HCl salt of 5,6-diethyl-indan-2-ylamine, a compound of formula XVII where R³, R⁴ and R⁷ are H, R⁵ and R⁶ are each CH₃CH₂-, R³⁰ is hydrogen and n is 1.

¹H-NMR (d₆-DMSO) ppm: 8.7 (3H, bd s, NH₃); 7.3 (2H, s, Ar); 4.2 (1H, bd s, CH); 3.5 (2H, dd, CH₂); 3.3 (2H, dd, CH₂); 2.8 (4H, q, CH₂CH₃); 1.4 (6H, t, CH₃).

Other compounds of formula XVII are prepared by procedures analogous to those used for Intermediate 1 or starting from available compounds and using procedures analogous to Preparations 3 and 4. These compounds of formula XVII are shown in the following table, R³ being hydrogen and n being 1 for all compounds.

Intermediate	R ⁴	R ⁵	R ⁶	R ⁷
2	CH ₃ CH ₂	H	H	CH ₃ CH ₂

3	H	-(CH ₂) ₄ -	H
4	H	-O(CH ₂) ₂ O-	H
5	H	CH ₃ (CH ₂) ₃ CH ₃ (CH ₂) ₃	H
6	H	CH ₃ (CH ₂) ₂ CH ₃ (CH ₂) ₂	H
7	H	CH ₃ O CH ₃ O	H

Intermediate 2 : ES + MS m/e (MH⁺) : 204

Intermediate 3 : ¹H-NMR (d6-DMSO) ppm: 8.1 (3H, bd s, NH₃); 6.9 (2H, s, Ar); 3.9 (1H, bd s, CH); 3.2 (2H, dd, CH₂); 2.8 (2H, dd, CH₂); 2.7 (4H, m, CH₂Ar); 1.7 (6H, t, CH₂).

Intermediate 4 : ¹H-NMR (d6-DMSO) ppm: 8.3 (3H, bds, NH₃); 6.85 (2H, s, Ar); 4.2 (4H, s, 2CH₂); 3.1 (2H, dd, CH₂); 2.85 (2H, dd, CH₂).

Intermediate 5 : ¹H-NMR (d6-DMSO) ppm: 6.9 (2H, s, Ar); 3.8 (1H, m, CH); 3.1 (2H, dd, CH₂); 2.6 (2H, dd, CH₂); 2.5 (4H, t, 2CH₂); 1.65 (2H, bds, NH₂); 1.55 (4H, m, 2CH₂); 1.4 (4H, m, 2CH₂); 0.95 (6H, t, 2CH₃).

Intermediate 6 : ¹H-NMR (d6-DMSO) ppm: 8.1 (3H, bd s, NH₃); 7.0 (2H, s, Ar); 3.9 (1H, bd s, CH); 3.2 (2H, dd, CH₂); 2.8 (2H, dd, CH₂); 2.5 (4H, q, EtCH₂Ar); 1.6 (4H, q, CH₂), 0.9 (6H, t, CH₃).

Intermediate 7 : ¹H-NMR (d6-DMSO) ppm: 8.3 (3H, bd s, NH₃), 6.9 (2H, s, H-Ar), 3.9 (1H, bd m, CHN), 3.7 (6H, s, CH₃O), 3.2 (2H, dd, CH₂), 2.9 (2H, dd, CH₂).

Intermediate 8 - 2-(Trifluoroacetyl-amino)-5,6-bis(methoxymethyl)indane

According to the procedure of Magnus *et.al* (*Tetrahed. Lett.*, 34, 23-26 (1993)) a solution of commercially available 1,4-dimethoxy-2-butyne (1.32 g, 11.5 mmol) in nitrogen-degassed ethanol is heated to 80°C with stirring under a nitrogen atmosphere.

Tris(triphenylphosphine)rhodium chloride (64 mg, 0.07 mmol) and a solution of 2,2,2-trifluoro-N-[1-(2-propynyl)-3-butynyl]-acetamide (470 mg, 2.32 mmol; prepared from literature procedure: Romero, Arthur G.; Leiby, Jeffrey A PCT Int. Appl. WO 9623760) in nitrogen-degassed ethanol (2 ml) are added in portions over 2 hours. The mixture is stirred under nitrogen at 80°C for a further 3 hours. The solvent is removed under vacuo and the residue is purified by flash chromatography on silica gel, eluting with hexane/ethyl acetate (2:1)

¹H-NMR (CDCl₃) ppm: 2.9 (2H, dd), 3.35 (2H, dd), 3.45 (6H, s), 4.57 (4H, s), 4.85 (1H, m), 6.4 (1H, br s), 7.30 (2H, s).

Intermediate 9 - 2-Amino-5,6-bis(methoxymethyl)indane

A solution of potassium hydroxide (150 mg, 2.60 mmol) in water (0.5 ml) is added to a solution of 2-(trifluoroacetyl-amino)-5,6-bis(methoxymethyl)indane (240 mg, 0.75 mmol) in methanol (3 mL) and the mixture is heated at reflux for 2.5 hours. The solvent was removed *in vacuo* and the residue is partitioned between aqueous sodium hydroxide (10 mL) and ethyl acetate (20 mL). The organic extract is dried (MgSO₄) and the solvent is removed *in vacuo* to leave the product as a dark oil.

¹H-NMR (CDCl₃) ppm: 2.60 (2H, dd), 3.10 (2H, dd), 3.33 (6H, s), 3.75 (1H, m), 4.42 (4H, s), 7.17 (2H, s).

Intermediate 10 - 8-Hydroxy-5-[(indan-2-ylamino)-acetyl]-1H-quinolin-2-one

5-(Chloroacetyl)-8-hydroxy-2(1H)-quinolinone (25 mg, 0.105 mmol) prepared from literature procedure (Yoshizaki, Shiro; Tanimura, Kaoru; Tamada, Shigeharu; Yabuuchi, Youichi; Nakagawa, Kazuyuki. J. Med. Chem. (1976), 19(9), 1138-42) is reacted neat with indan-2-ylamine (205 mg, 1.21 mmol) at 25 °C for 2 hours. The reaction mixture is purified by flash chromatography (silica, CH₂Cl₂/ methanol 9:1). ES+ MS *m/e* 335 (MH⁺).

Intermediate 11

This compound of formula XVIII where Ar is a group of formula III, R²⁷, R²⁸ and R²⁹ are hydrogen, R², R³, R⁴ and R⁷ are hydrogen, and R⁵ and R⁶ are each methoxy, is prepared by a procedure analogous to that used for preparation of Intermediate 10. ES+MS *m/e*(MH⁺):395.

Intermediate 12 - 8-Benzyloxy-3-methyl-5-oxiranyl-1H-quinolin-2-one

8-Hydroxy-3-methyl-1H-quinolin-2-one is prepared according to the procedure of Wang *et al* (T.-C. Wang, Y.-L. Chen, K.-H. Lee, C.-C. Izeng *Synthesis* 1997, 87-90.).

¹H-NMR (d₄-CH₃OH) ppm: 2.14 (s, 3H), 6.84-6.89 (m, 1H), 6.95-7.03 (m, 2H), 6.90 (s, 1H), 7.71 (s, 1H).

8-Benzyloxy-3-methyl-1H-quinolin-2-one

Benzyl bromide (1.28 mL) is added to a suspension of potassium carbonate (2.98 g) in a solution of 8-hydroxy-3-methyl-1H-quinolin-2-one (1.26 g) in acetone (36 mL) at room temperature. The reaction mixture is refluxed for 18 hours, filtered, evaporated and purified by flash column chromatography on silica gel, eluting with 2 % methanol in dichloromethane.

¹H-NMR (CDCl₃) ppm: 2.11 (s, 3H), 5.13 (s, 2H), 6.92-6.98 (m, 1H), 7.02-7.08 (m, 2H), 7.29-7.40 (m, 5H), 7.57 (s, 1H), 9.23 (s, 1H).

8-Benzyloxy-5-bromo-3-methyl-1H-quinolin-2-one

A solution of bromine (0.57 g) in acetic acid (2 mL) is added dropwise to a solution of 8-benzyloxy-3-methyl-1H-quinolin-2-one (0.94g) and sodium acetate (0.96 g) in acetic acid (12 mL) at room temperature. The reaction mixture is stirred at room temperature for 3 hours, evaporated, the residue partitioned between water (5 mL) and ethyl acetate (5 mL), extracting a further 2x with ethyl acetate (5 mL). Combined organic extracts are dried over magnesium sulphate and purified by flash column chromatography on silica gel, eluting with 2% methanol in dichloromethane.

¹H-NMR (CDCl₃) ppm: 2.27 (s, 3H), 5.18 (s, 2H), 6.83 (d, 1H), 7.39 (d, 1H), 7.37-7.41 (m, 5H), 7.91 (s, 1H), 9.08 (s, 1H).

8-Benzyloxy-3-methyl-5-vinyl-1H-quinolin-2-one

Palladium terakis(triphenylphosphine) (30 mg) is added to a solution of 8-benzyloxy-5-bromo-3-methyl-1H-quinolin-2-one (239 mg) and tributylvinyltin (203 µL) in toluene (7 mL) at room temperature. The reaction mixture is heated for 2 hours at 100 °C, cooled to room temperature, evaporated and the product purified by flash column chromatography on silica gel, eluting with 2% ethyl acetate in dichloromethane.

¹H-NMR (CDCl₃) ppm: 2.24 (s, 3H), 5.18 (s, 2H), 5.32-5.39 (m, 1H), 5.61-5.68 (m, 1H), 6.95 (d, 1H), 7.09-7.20 (m, 1H), 7.21-7.26 (m, 2H), 7.31-7.43 (m, 4H), 7.89 (s, 1H), 9.20 (s, 1H).

8-Benzyloxy-3-methyl-5-oxiranyl-1H-quinolin-2-one

To 8-benzyloxy-3-methyl-5-vinyl-1H-quinolin-2-one (300 mg) is added to a 0.1M solution of dimethyldioxirane in acetone (12.4 mL). After stirring at room temperature for 2 hours, the solvent is removed *in vacuo* to yield the product.

¹H-NMR (CDCl₃) ppm: 2.23 (s, 3H), 2.77-2.81 (m, 1H), 3.18-3.23 (m, 1H), 4.17-4.21 (m, 1H), 5.18 (s, 2H), 6.91 (d, 1H), 7.01 (d, 1H), 7.93 (s, 1H), 9.10 (s, 1H).

Intermediate 13 - 8-Benzyloxy-5-[2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-3-methyl-1H-quinolin-2-one

A solution of Intermediate 12 (65 mg) and 5,6-diethyl-indan-2-ylamine (120 mg) in DMSO (1.5 mL) is heated for 18 hours at 90 °C. The solvent is removed *in vacuo*, and the product purified by flash chromatography on silica gel, eluting with 10% methanol in dichloromethane.

¹³C-NMR (d4-CH₃OH) ppm: 15.96, 17.14, 26.33, 36.77, 53.34, 59.82, 67.33, 71.73, 112.09, 118.98, 121.73, 125.42, 128.74, 129.24, 129.47, 129.61, 131.84, 134.56, 137.52, 137.64, 142.29, 145.94, 164.02.

Intermediate 14 - 8-Methoxymethoxy-6-methyl-5-oxiranyl-1H-quinolin-2-one

8-Hydroxy-6-methyl-1H-quinolin-2-one is prepared according to the procedure of Wang *et al* (T.-C. Wang, Y.-L. Chen, K.-H. Lee, C.-C. Izeng *Synthesis* **1997**, 87-90.).

¹H-NMR (d6-DMSO) ppm: 2.26 (s, 3H), 6.45 (d, 1H), 6.79 (s, 1H), 6.90 (s, 1H), 7.78 (d, 1H).

5-Bromo-8-hydroxy-6-methyl-1H-quinolin-2-one

A 45 % solution of hydrobromic acid in acetic acid (324 µL) is added dropwise to a solution of 8-hydroxy-6-methyl-1H-quinolin-2-one (316 mg) in dimethylsulphoxide (9 mL) at room temperature. The reaction mixture is allowed to stand for 18 hours at room temperature and the solvent removed *in vacuo*.

¹H-NMR (d6-DMSO) ppm: 2.33 (s, 3H), 6.58 (d, 1H), 6.92 (s, 1H), 8.03 (d, 1H), 10.44 (s, 1H), 10.67 (s, br, 1H).

5-Bromo-8-methoxymethoxy-6-methyl-1H-quinolin-2-one

Methoxymethyl chloride (410 µL) was added to a suspension of potassium carbonate (1.24 g) in a solution of 5-bromo-8-hydroxy-6-methyl-1H-quinolin-2-one (480 mg) in dimethylformamide (9 mL) at 0 °C. The reaction mixture is stirred for 18 hours at room temperature, filtered, the solvent removed *in vacuo*, and the product purified by flash column chromatography on silica gel, eluting with 2 % methanol in dichloromethane.

¹³C-NMR (CDCl₃) ppm: 23.42, 56.52, 95.07, 115.78, 116.19, 119.32, 123.30, 128.13, 132.14, 139.78, 141.78, 161.32.

8-Methoxymethoxy-6-methyl-5-vinyl-1 H-quinolin-2-one

Bis-(triphenylphosphine)palladium (II) chloride (98 mg) is added to a solution of 5-bromo-8-methoxymethoxy-6-methyl-1H-quinolin-2-one (410 mg) and tributylvinyltin (603 µL) in dimethylformamide (14 mL) at room temperature. The reaction mixture is heated for 24 hours at 90 °C, evaporated and purified by flash column chromatography on silica gel, eluting with 2 % methanol in dichloromethane.

¹H-NMR (CDCl₃) ppm: 2.19 (s, 3H), 3.41 (s, 3H), 5.18 (d, 1H), 5.20 (s, 2H), 5.60 (d, 1H), 6.52 (d, 1H), 6.63-6.69 (m, 1H), 6.96 (s, 1H), 7.95 (d, 1H), 9.78 (s, 1H).

8-Methoxymethoxy-6-methyl-5-oxiranyl-1H-quinolin-2-one is obtained from 8-methoxymethoxy-6-methyl-5-vinyl-1H-quinolin-2-one

(186 mg) according to the last step of the procedure for Intermediate 12.

¹H-NMR (CDCl₃) ppm: 2.38 (s, 3H), 2.68-2.72 (m, 1H), 3.19-3.23 (m, 1H), 3.43 (s, 3H), 3.97-4.01 (m, 1H), 5.21 (s, 2H), 6.60 (d, 1H), 6.98 (s, 1H), 8.22 (d, 1H), 9.09 (s, 1H).

Intermediate 15, (R)-2-(4-benzyloxy-3-nitrophenyl)-oxirane, is prepared according to the procedure of R. Hett et al, Tetrahedron Lett. (1997), 38(7), 1125-1128.

Intermediate 16 - (S)-8-Benzyloxy-5-oxiranyl-1H-quinolin-2-one

8-Benzyloxy-5-((S)-2-chloro-1-hydroxy-ethyl)-1H-quinolin-2-one

(S)-2-methyl-CBS-oxazaborolidine, 1M in toluene (0.30mL, 0.30mmol) is added to dry THF (tetrahydrofuran) (10mL) in an oven dried flask. Borane-THF complex, 1M in THF (3.05mL) is then added dropwise and the solution is stirred at room temperature for 15 minutes and then cooled to 0°C. 8-Benzyloxy-5-chloroacetyl-1H-quinolin-2-one (1.00g), prepared as described in WO95/25104, is then added in small portions over a period of 30 minutes. The reaction mixture is stirred at 0°C. The reaction is shown to be complete by TLC (thin layer chromatography) after 15 minutes. The reaction mixture is quenched with methanol (1mL), the solvent is removed *in vacuo* and the residue is partitioned between 0.2M H₂SO₄ (100mL) and CHCl₃ (100mL). The organic layer is dried over MgSO₄, filtered and the solvent is removed *in vacuo*. Crystallised from ethyl acetate. TLC (silica, dichloromethane / methanol 25:1 R_f = 0.30).

(S)-8-Benzyloxy-5-oxiranyl-1H-quinolin-2-one

8-Benzyloxy-5-((S)-2-chloro-1-hydroxy-ethyl)-1H-quinolin-2-one (0.55g) is dissolved in acetone (20mL). K₂CO₃ (0.58g) is added and the reaction mixture is refluxed. The reaction is shown to be complete by TLC after 18 hours. The solvent is removed *in vacuo* and the residue is partitioned between ethyl acetate (100mL) and water (100mL). The organic layer is dried over MgSO₄, filtered and the solvent is removed *in vacuo*. The product is triturated with diethyl ether, filtered and dried. TLC (silica, dichloromethane / methanol 25:1 R_f = 0.45).

Intermediate 17 - 6,7,8,9-Tetrahydro-5H-benzocyclohepten-7-ylamine**Benzyl-(6,7,8,9-tetrahydro-5H-benzocyclohepten-7-yl)-amine**

5,6,8,9-Tetrahydro-benzocyclohepten-7-one (3.00g) and benzylamine (2.00g) are dissolved in ethanol (50mL). A catalytic amount of 10% palladium on charcoal is added and the reaction mixture is placed under an atmosphere of hydrogen. The reaction mixture is stirred at r.t. The reaction is shown to be complete by TLC after 24 hours. The catalyst is filtered off and the solvent is removed *in vacuo*. The product is not purified further. TLC (silica, n-hexane /ethyl acetate 1:2 R_f = 0.50).

6,7,8,9-Tetrahydro-5H-benzocyclohepten-7-ylamine

Benzyl-(6,7,8,9-tetrahydro-5H-benzocyclohepten-7-yl)-amine (2.80g) is dissolved in methanol (100mL) and the compound is deprotected by adding a catalytic amount of 10% palladium on charcoal and placing the solution under an atmosphere of hydrogen. The reaction is shown to be complete by TLC after 24 hours. The catalyst is filtered off and the solvent is removed *in vacuo*. The product is not purified further. TLC (silica, dichloromethane / methanol 25:1 R_f = 0.15).

Intermediate 18 - Benzyl-(5,6-diethyl-indan-2-yl)-amine**N-(5,6-Diethyl-indan-2-yl)-benzamide**

5,6-Diethyl-indan-2-ylamine (4.10g) is dissolved in dichloromethane (DCM) (150mL) and triethylamine (2.41g) is added. Benzoyl chloride (3.20g) is then added dropwise and the reaction mixture is stirred at room temperature. The reaction is shown to be complete by TLC after 1 hour. The solution is washed with 0.2M HCl (100mL), water (100mL) and brine (100mL). The organic layer is dried over $MgSO_4$, filtered and the solvent is removed *in vacuo*. Crystallised from ethyl acetate. TLC (silica, dichloromethane / methanol 10:1 R_f = 0.85).

Benzyl-(5,6-diethyl-indan-2-yl)-amine

N-(5,6-Diethyl-indan-2-yl)-benzamide (3.30g) is dissolved in dry THF (100mL). Lithium aluminium hydride, 1M in THF (22.52ml) is then added dropwise. The reaction mixture is stirred at 50°C. The reaction is shown to be complete by TLC after 6 hours. The reaction mixture is allowed to cool, poured slowly into ice-water (200mL) and extracted with diethyl ether (2 x 150mL). The organic layer is dried over $MgSO_4$, filtered and the solvent is

removed *in vacuo*. The product is not purified further. TLC (silica, n-hexane / ethyl acetate 2:1 R_f = 0.20).

Intermediate 19 - (R)-1-(3-Amino-4-benzyloxy-phenyl)-2-[benzyl-(5,6-diethyl-indan-2-yl)-amino]-ethanol

(R)-2-[Benzyl-(5,6-diethyl-indan-2-yl)-amino]-1-(4-benzyloxy-3-nitro-phenyl)-ethanol

The title compound is prepared from Intermediate 15 (3.01g) and Intermediate 18 (3.10g) by an analogous procedure to that used to prepare (S)-8-Benzyloxy-5-[2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-1H-quinolin-2-one in Example 19. The reaction is shown to be complete by TLC after 24 hours. The product is purified by flash column chromatography (silica, n-hexane / ethyl acetate 4:1). TLC (silica, n-hexane / ethyl acetate 4:1 R_f = 0.25).

(R)-1-(3-Amino-4-benzyloxy-phenyl)-2-[benzyl-(5,6-diethyl-indan-2-yl)-amino]-ethanol

(R)-2-[Benzyl-(5,6-diethyl-indan-2-yl)-amino]-1-(4-benzyloxy-3-nitro-phenyl)-ethanol (3.00g) is dissolved in THF (50mL) and toluene (50mL). A catalytic amount of PtO_2 is added and the solution is stirred under an atmosphere of H_2 . The reaction is shown to be complete by TLC after 6 hours. The catalyst is filtered off and the solvent is removed *in vacuo*. The product is not purified further. TLC (silica, n-hexane / ethyl acetate 1:1 R_f = 0.75).

Intermediate 20 - 1-(3-Amino-4-benzyloxy-phenyl)-2-[benzyl-(5,6-diethyl-indan-2-yl)-amino]-ethanone

2-[Benzyl-(5,6-diethyl-indan-2-yl)-amino]-1-(4-benzyloxy-3-nitro-phenyl)-ethanone

1-(4-Benzyloxy-3-nitro-phenyl)-2-bromo-ethanone (2.00g) (Prepared following procedure; Hett, Robert; Fang, Qun Kevin; Gao, Yun; Hong, Yaping; Butler, Hal T.; Nie, Xiaoyi; Wald, Stephen A. *Tetrahedron Lett.* 1997, 38, 1125-1128.) is dissolved in methymethylketone (100mL). Triethylamine (0.64g) is added followed by benzyl-(5,6-diethyl-indan-2-yl)-amine (1.60g). The reaction mixture is then refluxed. The reaction is shown to be complete by TLC after 3 hours. The solvent is removed *in vacuo* and the product is purified by flash column chromatography (silica, n-hexane / ethyl acetate 4:1). TLC (silica, n-hexane / ethyl acetate 2:1 R_f = 0.75).

1-(3-Amino-4-benzyloxy-phenyl)-2-[benzyl-(5,6-diethyl-indan-2-yl)-amino]-ethanone is prepared from 2-[benzyl-(5,6-diethyl-indan-2-yl)-amino]-1-(4-benzyloxy-3-nitro-phenyl)-

ethanone (1.50g) by an analogous procedure to that used to prepare (R)-1-(3-Amino-4-benzyloxy-phenyl)-2-[benzyl-(5,6-diethyl-indan-2-yl)-amino]-ethanol in Example 19. The reaction is shown to be complete by TLC after 48 hours. The catalyst is filtered off and the solvent is removed *in vacuo*. The product is purified by flash column chromatography (silica, n-hexane / ethyl acetate 4:1). TLC (silica, n-hexane / ethyl acetate 2:1 R_f = 0.70). ^1H NMR [CDCl_3 , 400MHz] δ 1.20 (6H, t), 1.60 (2H, broad), 2.60 (4H, q), 3.00 (4H, m), 3.90 (6H, m), 5.15 (2H, s), 6.80 (1H, d), 6.95 (2H, s), 7.30 (12H, m).

Intermediate 21 - Benzyl-(4,5,6,7-tetramethyl-indan-2-yl)-amine

3-Chloro-1-(2,3,4,5-tetramethyl-phenyl)-propan-1-one is prepared from 1,2,3,4-tetramethylbenzene and 3-chloro propionyl chloride by a procedure analogous to that of Preparation 1. ^1H NMR (CD_3OD) ppm: 7.5 (1H, s); 4.2 (2H, t); 3.6 (2H, t); 2.6 (3H, s); 2.57 (3H, s); 2.52 (3H, s); 2.5 (3H, s).

4,5,6,7-Tetramethyl-indan-1-one is prepared from 3-chloro-1-(2,3,4,5-tetramethyl-phenyl)-propan-1-one by a procedure analogous to that of Preparation 2. ^1H NMR (CD_3OD) ppm: 3.2 (2H, t); 2.9 (2H, t); 2.85 (3H, s); 2.6 (3H, s); 2.55 (3H, s); 2.5 (3H, s).

4,5,6,7-Tetramethyl-indan-1,2-dione 2-oxime is prepared from 4,5,6,7-tetramethyl-indan-1-one by a procedure analogous to that of Preparation 3. ^1H NMR ($\text{d}_6\text{-DMSO}$) ppm: 12.4 (1H, s); 3.65 (2H, s); 2.7 (3H, s); 2.4 (3H, s); 2.3 (6H, s).

2-Amino-4,5,6,7-tetramethyl-indan-1-one hydrochloride is prepared from 4,5,6,7-tetramethyl-indan-1,2-dione 2-oxime by a procedure analogous to that of Preparation 4. ^1H NMR ($\text{d}_6\text{-DMSO}$) ppm: 9.0 (3H, bd s); 4.5 (1H, bd t); 3.7 (1H, dd); 3.2 (1H, dd); 2.8 (3H, s); 2.6 (3H, s); 2.5 (6H, 2 s).

N-(4,5,6,7-Tetramethyl-1-oxo-indan-2-yl)-benzamide

Benzoyl chloride (1.635g) is added dropwise to 4,5,6,7-tetramethyl-indan-1,2-dione 2-oxime (2.53g) and triethylamine (2.25g) in anhydrous dichloromethane (60ml) at 0°C. The reaction mixture is stirred at room temperature for 1.5 hours after which the solid product is filtered off and allowed to stir with water (150ml), refiltered and dried. The organic filtrate is washed with 1M HCl, 10% brine, saturated sodium bicarbonate solution, 10% brine and

treated with magnesium sulphate. After filtration, the solvent is removed in vacuo and the product triterated with diethyl ether, filtered and dried.

¹H NMR (CDCl₃) ppm: 7.8 (2H, d); 7.45 (1H, m), 7.4 (2H, m); 6.8 (1H, bd d); 4.6 (1H, m); 3.8 (1H, dd); 2.8 (1H, dd); 2.55 (3H, s); 2.25 (3H, s); 2.15 (6H, 2 s).

N-(1-Hydroxy-4,5,6,7-tetramethyl-indan-2-yl)-benzamide

Sodium borohydride (213 mg) is added to N-(4,5,6,7-tetramethyl-1-oxo-indan-2-yl)-benzamide (495 mg) in chloroform (20 ml) and methanol (20 ml). The reaction mixture is stirred at room temperature for 2 hours, drowned with water (50 ml) and chloroform (20 ml) added. The aqueous phase is washed with chloroform (x2) and the organic layers combined, treated with magnesium sulphate, filtered and the solvent removed in vacuo.

¹H NMR (CDCl₃) ppm: 7.65 (2H, d); 7.4 (1H, m), 7.35 (2H, m); 6.3 (1H, bd d); 5.15 (1H, d); 4.5 (1H, m); 3.7 (1H, bd s); 3.5 (1H, dd); 2.65 (1H, dd); 2.25 (3H, s); 2.15 (9H, 3 s).

N-(4,5,6,7-Tetramethyl-indan-2-yl)-benzamide is prepared from N-(1-Hydroxy-4,5,6,7-tetramethyl-indan-2-yl)-benzamide by a procedure analogous to that of Preparation 4.

¹H NMR (CDCl₃) ppm: 7.65 (2H, d); 7.4 (1H, m), 7.3 (2H, m); 6.25 (1H, bd d); 4.85 (1H, m); 3.35 (1H, dd); 2.80 (1H, dd); 2.1 (12H, 2s).

Benzyl-(4,5,6,7-tetramethyl-indan-2-yl)-amine

1M Lithium aluminium hydride (2.4 ml) in tetrahydrofuran is added dropwise to a solution of N-(4,5,6,7-tetramethyl-indan-2-yl)-benzamide (352 mg) in anhydrous THF (10ml) under nitrogen at room temperature. The reaction mixture is allowed to stir at 50°C for 20 hours. After 4 hours more 1M Lithium aluminium hydride (1.2 ml, 1.20 mmole) in THF is added. On cooling the reaction mixture is quenched with ice water. The aqueous phase is washed with diethyl ether (x3) and the organic layers combined, treated with magnesium sulphate, filtered and the solvent removed in vacuo.

¹H NMR (CDCl₃) ppm: 7.25 (4H, m); 7.15 (1H, m); 3.8 (2H, s); 3.55 (1H, m); 3.1 (2H, dd); 2.7 (2H, dd); 2.1 (12H, 2s).

Intermediate 22 - Benzyl-(2,3,5,6,7,8-hexahydro-1H-cyclopenta[b]naphthalen-2-yl)-amine

According to the procedure of A.F.Abdel-Magid, et.al. *J.Org.Chem.* **1996**, 61, 3849-3862. triethylamine (0.87 mL, 6.17 mmol) is added to a stirred suspension 2,3,5,6,7,8-hexahydro-1H-cyclopenta[b]naphthalen-2-ylamine in 1,2-dichloroethane (30 mL) under nitrogen at room temperature. Benzaldehyde (0.52 mL, 5.14 mmol) is then added followed by sodium

triacetoxyborohydride (1.64 g, 7.7 mmol) and acetic acid (0.44 mL, 7.7 mmol). The reaction is stirred at room temperature for 18 hours. After diluting with dichloromethane the mixture is washed with aqueous NaOH (50 mL, 1M) followed by brine. Removal of the solvent and chromatography (silica, ethyl acetate / hexane, 2:1) affords an oil.

¹H-NMR (CDCl₃) ppm: 1.70 (m, 4H), 2.65 (m, 4H), 2.68 (dd, 2H), 3.05 (dd, 2H), 3.58 (m, 1H), 3.78 (s, 2H), 6.83 (s, 2H), 7.25 (m, 5H).

Intermediate 23 - 2-Methyl-indan-2-ylamine

2-Amino-2-methyl-indan-1-one

According to the procedure of Farnum *et.al* (*Synthesis* 1972, 191-192.), water (1.35 L) is stirred at 80°C and de-gassed by periodic evacuating and flushing with nitrogen (3 x). K₃FeCN₆ (202 g, 615 mmol) and 2-methyl-indan-1-one (20 g, 137 mmol) are added. The mixture is stirred rapidly under nitrogen at 80°C while aqueous concentrated ammonia solution (105 mL) is added over 30 minutes. Stirring is continued at 80°C for 20 hours. When cool, the solution is made alkaline by addition of sodium hydroxide (2 g) and extracted with ethyl acetate (2 x 200 mL). The organic extract is concentrated to a volume of 200 ml and the product is extracted into aqueous HCl (200 mL, 1M). The acidic aqueous phase is separated, basified with sodium hydroxide, and extracted with ethyl acetate (2 x 100 mL). The organic layer is separated, dried (Na₂SO₄) and the solvent removed to give an orange oil.

¹H-NMR (CDCl₃) ppm: 1.38 (s, 3H), 1.8 (br. s, 2H), 3.07 (d, 1H), 3.25 (d, 1H), 3.45 (m, 2H), 7.65 (t, 1H), 7.80 (d, 1H).

2,2,2-Trifluoro-N-(2-methyl-1-oxo-indan-2-yl)-acetamide

2-Amino-2-methyl-indan-1-one (16.4 g) in THF (100 mL) is cooled to 0°C under nitrogen. Triethylamine (21 ml) is added followed by slow addition of trifluoroacetic anhydride (18.5 ml). The reaction is stirred at room temperature overnight then the solvents are removed. The residue is dissolved in dichloromethane and washed with aqueous HCl followed by aqueous NaOH. The organic extract is dried (MgSO₄) and the solvent is removed. The product is purified by chromatography (silica, ethyl acetate) to give a cream solid.

¹H-NMR (CDCl₃) ppm: 1.52 (s, 3H), 3.44 (d, 1H), 3.55 (d, 1H), 7.05 (br.s, 1H), 7.43 (m, 2H), 7.70 (t, 1H), 7.87 (d, 1H).

2,2,2-Trifluoro-N-(2-methyl-indan-2-yl)-acetamide

2,2,2-Trifluoro-N.-(2-methyl-1-oxo-indan-2-yl)-acetamide (3.41 g) in acetic acid (25 mL) and H₂SO₄ (0.5 mL) is stirred under hydrogen in the presence of 10% Pd/C at room temperature for 18 hours. The mixture is filtered through celite and the filtrate is concentrated *in vacuo*. After diluting with water the mixture is extracted with diethyl ether. The organic phase is removed, washed several times with aqueous sodium bicarbonate and dried (Na₂SO₄). The solvent is removed to give an oil which solidifies.

¹H-NMR (CDCl₃) ppm: 1.55 (s, 3H), 3.05 (d, 2H), 3.28 (d, 2H), 6.28 (br.s, 1H), 7.12 (s, 4H).

2-Methyl-indan-2-ylamine

A stirred solution of 2,2,2-trifluoro-N.-(2-methyl-indan-2-yl)-acetamide (6.70 g) and NaOH (4.0 g) in methanol (100 mL) and water (1 mL) is heated at 70°C for 2 hours. The solvent is removed and the residue is partitioned between aqueous HCl (100 mL, 2M) and ethyl acetate (100 mL). The aqueous extract is separated, basified with aq. NaOH, and extracted with ethyl acetate. The organic phase is separated, dried (MgSO₄), and the solvent is removed to give an orange oil which solidifies.

¹H-NMR (CDCl₃) ppm: 1.19 (s, 3H), 1.5 (br.s, 2H), 2.65 (d, 2H), 2.79 (d, 2H), 6.97 (m, 4H).

Intermediate 24 - 2-Methyl-2,3,5,6,7,8-hexahydro-1H-cyclopenta[b]naphthalen-2-ylamine

1-(5,6,7,8-Tetrahydro-naphthalen-2-yl)-propan-1-one

Propionyl chloride (17.5 mL) and 1,2,3,3-tetrahydronaphthalene (27.5 mL) are added slowly over 1 hour to a stirred solution of AlCl₃ (61.3 g) in nitromethane (200 mL) at 0°C. After stirring at room temperature for 18 hours the reaction is cautiously added to a mixture of ice and concentrated HCl. The product is extracted with ethyl acetate, washed with brine and dried (Na₂SO₄).

¹H-NMR (CDCl₃) ppm: 1.15 (t, 3H), 1.72 (m, 4H), 2.72 (m, 4H), 2.88 (q, 2H), 7.04 (d, 1H), 7.60 (m, 1H).

2-Methyl-2,3,5,6,7,8-hexahydro-cyclopenta[b]naphthalen-1-one

According to the procedure of Bhattacharya *et.al* (*Synth. Commun* 1996., 26, 1775-1784.) a mixture of 1-(5,6,7,8-tetrahydro-naphthalen-2-yl)-propan-1-one (37.6 g), hexamethylenetetramine (44.9 g) and acetic anhydride (38.8 mL) is heated with stirring at 80°C for 23 hours. The mixture is allowed to cool, and added slowly to a stirred mixture of ethyl acetate (200 mL) and aqueous sodium hydroxide (200 mL, 2M). The organic layer

is separated, washed with aqueous HCl, brine, and dried (Na_2SO_4). The solvent is removed to give a brown oil. This is added cautiously to concentrated sulfuric acid (120 mL) and the resulting mixture is heated at 55°C for 5 hours followed by room temperature for 18 hours. The reaction is diluted with water and extracted with dichloromethane. After drying (Na_2SO_4) the solvent is removed to give an oil. The product is purified by chromatography (silica, ethyl acetate / hexane) to give a geometrical mixture of isomers containing 2-Methyl-1,2,6,7,8,9-hexahydro-cyclopenta[a.]naphthalen-3-one and the title compound.

$^1\text{H-NMR}$ (CDCl_3) ppm (mixture): 1.4 (m, 3H), 1.9 (m, 4H), 2.5-3.0 (m, 6H), 3.35 (m, 1H), 7.15 (m, 1H), 7.55 (m, 1H).

2,2,2-Trifluoro-N-(2-methyl-1-oxo-2,3,5,6,7,8-hexahydro-1H-cyclopenta[b]naphthalen-2-yl)-acetamide

This compound is prepared from an isomeric mixture, containing 2-methyl-2,3,5,6,7,8-hexahydro-cyclopenta[b]naphthalen-1-one, according to the procedure used for the preparation of 2,2,2-trifluoro-N-(2-methyl-1-oxo-indan-2-yl)-acetamide. The isomeric mixture of products is recrystallised from ethyl acetate/hexane to give a 4:1 mixture in favour of the title compound.

$^1\text{H-NMR}$ (CDCl_3) ppm (major component): 1.55(s, 3H), 1.85 (m, 4H), 2.87 (m, 4H), 6.88 (br.s, 1H), 7.18 (s, 1H), 7.57 (s, 1H).

TOF MS ES⁻ *m/e* 310 ($\text{M} - \text{H}^-$)

2-Methyl-2,3,5,6,7,8-hexahydro-1H-cyclopenta[b]naphthalen-2-ylamine

A 4:1 mixture of geometrical isomers, containing predominantly 2,2,2-trifluoro-N-(2-methyl-1-oxo-2,3,5,6,7,8-hexahydro-1H-cyclopenta[b]naphthalen-2-yl)-acetamide, is hydrogenated over Pd/C in acetic acid/ H_2SO_4 and the products are saponified with NaOH according to the procedures described for the preparation of 2-methyl-indan-2-ylamine. The resulting product mixture is recrystallised repeatedly from hexane to give the title compound, a single isomer.

$^1\text{H-NMR}$ (CDCl_3) ppm: 1.40 (s, 3H), 1.6 (br.s, NH_2), 1.75 (m, 4H), 3.75 (m, 4H), 2.78 (d, 2H), 2.94 (d, 2H), 6.93 (s, 2H).

Intermediate 25 - 2-Ethyl-indan-2-ylamine

2-Ethyl-indan-1-one is prepared from benzene following analogous procedures to those used for 2-methyl-2,3,5,6,7,8-hexahydro-cyclopenta[b]naphthalen-1-one.

¹H-NMR (CDCl₃) ppm: 0.97 (t, 3H), 1.50 (m, 1H), 1.90 (m, 1H), 2.55 (m, 1H), 2.75 (dd, 1H), 3.25 (q, 1H), 7.29 (t, 1H), 7.39 (d, 1H), 7.50 (t, 1H), 7.69 (d, 1H).

2-Ethyl-indan-2-ylamine is prepared from 2-ethyl-indan-1-one by procedures analogous to those used for Intermediate 23.

¹H-NMR (CDCl₃) ppm: 1.05 (t, 3H), 1.5 (br.s, NH₂), 2.70 (q, 2H), 2.75 (d, 2H), 3.01 (d, 2H), 7.20 (m, 4H).

Intermediate 26 - 2,5,6-Trimethyl-indan-2-ylamine

2,5,6-Trimethyl-indan-2-ylamine is prepared from 1,2-dimethylbenzene by procedures analogous to those used for 2-methyl-2,3,5,6,7,8-hexahydro-1H-cyclopenta[b]naphthalen-2-ylamine.

¹H-NMR (CDCl₃) ppm: 1.29 (s, 3H), 2.16 (s, 6H), 2.69 (d, 2H), 2.84 (d, 2H), 2.89 (s, 2H).

Intermediate 27 - Acetic acid (R)-1-(3-amino-4-benzyloxy-phenyl)-2-[benzyl-(2-methyl-indan-2-yl)-amino]-ethyl ester

(R)-2-[Benzyl-(2-methyl-indan-2-yl)-amino]-1-(4-benzyloxy-3-nitro-phenyl)-ethanol

The title compound is prepared from (R)-2-(4-benzyloxy-3-nitro-phenyl)-oxirane (2.52g) and benzyl-(2-methyl-indan-2-yl)-amine (2.20g) by an analogous procedure to that used to prepare (S)-8-Benzyloxy-5-[2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-1H-quinolin-2-one in Example 19. The reaction is shown to be complete by TLC after 24 hours. The product is purified by flash column chromatography (silica, n-hexane / ethyl acetate 4:1). TLC (silica, n-hexane / ethyl acetate 4:1 R_f = 0.30).

¹H NMR [CDCl₃, 400MHz] d 1.20 (3H, s), 2.65 (1H, m), 2.75 (1H, m), 2.90 (2H, m), 3.25 (2H, m), 3.60 (1H, d), 3.70 (1H, broad), 3.80 (1H, d of d), 4.10 (1H, d), 5.20 (2H, s), 7.00 (1H, d), 7.20 (4H, m), 7.35 (11H, m), 7.60 (1H, d).

Acetic acid (R)-2-[benzyl-(2-methyl-indan-2-yl)-amino]-1-(4-benzyloxy-3-nitro-phenyl)-ethyl ester

(R)-2-[Benzyl-(2-methyl-indan-2-yl)-amino]-1-(4-benzyloxy-3-nitro-phenyl)-ethanol (2.75g) is dissolved in pyridine (15mL). Acetic anhydride (1.66g) is added and the reaction mixture is stirred at room temperature. The reaction is shown to be complete by TLC after 18 hours. Water (10mL) is added to quench the reaction. Ethyl acetate(250mL) is added and the

solution is washed with 1M KHSO₄ (3 x 100mL), saturated NaHCO₃ (100mL), water (100mL) and brine (100mL). The organic layer is dried over MgSO₄, filtered and the solvent is removed *in vacuo*. The product is not purified further. TLC (silica, n-hexane / ethyl acetate 4:1 R_f = 0.40).

¹H NMR [CDCl₃, 400MHz] d 1.20 (3H, s), 1.90 (3H, s), 2.80 (3H, m), 3.00 (1H, d), 3.10 (1H, m), 3.20 (1H, d), 3.75 (1H, d), 3.90 (1H, d), 5.20 (2H, s), 5.25 (1H, m), 6.95 (1H, d), 7.10 (4H, m), 7.30 (11H, m), 7.55 (1H, d).

Acetic acid (R)-1-(3-amino-4-benzyloxy-phenyl)-2-[benzyl-(2-methyl-indan-2-yl)-amino]-ethyl ester

The title compound is prepared from acetic acid (R)-2-[benzyl-(2-methyl-indan-2-yl)-amino]-1-(4-benzyloxy-3-nitro-phenyl)-ethyl ester (2.90g) by an analogous procedure to that used to prepare (R)-1-(3-amino-4-benzyloxy-phenyl)-2-[benzyl-(5,6-diethyl-indan-2-yl)-amino]-ethanol in Example 19. The reaction is shown to be complete by TLC after 6 hours. The catalyst is filtered off and the solvent is removed *in vacuo*. The product is not purified further. TLC (silica, n-hexane / ethyl acetate 2:1 R_f = 0.60).

¹H NMR [CDCl₃, 400MHz] d 1.10 (3H, s), 1.80 (3H, s), 2.70 (3H, m), 3.05 (2H, m), 3.15 (1H, d), 3.65 (2H, broad), 3.75 (1H, d), 3.90 (1H, d), 4.95 (2H, s), 5.20 (1H, m), 6.40 (2H, m), 6.65 (1H, d), 7.20 (14H, m).

Intermediate 28 – Benzyl-(2,5,6-trimethyl-indan-2-yl)-amine

N-(2,5,6-Trimethyl-indan-2-yl)-benzamide

Intermediate 26 is treated with benzoyl chloride in dichloromethane / triethylamine for 1 hour. The mixture is washed with 1N HCl, then with saturated NaHCO₃ solution, dried (Na₂SO₄) and evaporated. The residue is triturated with ether / hexane to give white crystals. ¹H-NMR (CDCl₃) ppm: 1.60 (s, 3H), 2.18 (s, 6H), 3.02 (d, 2H), 3.30 (d, 2H), 6.17 (br.s, NH), 6.90 (s, 2H), 7.34 (m, 2H), 7.40 (m, 1H), 7.63 (d, 1H).

Benzyl-(2,5,6-trimethyl-indan-2-yl)-amine

To a solution of N-(2,5,6-trimethyl-indan-2-yl)-benzamide in THF under nitrogen is added LiAlH₄ and the mixture refluxed for 48 hours. Quenched at 0°C with ice / water and extracted with ether, dried (Na₂SO₄) and solvent removed *in vacuo*. Purification by chromatography (silica, ethyl acetate / hexane 1:4) gives a colourless oil.

¹H-NMR (CDCl₃) ppm: 1.58 (s, 3H), 1.79 (br.s., NH), 2.40 (s, 6H), 3.00 (d, 2H), 3.20 (d, 2H), 3.99 (s, 2H), 7.15 (s, 2H), 7.37 - 7.53 (m, 5H).

Intermediate 29 – Acetic acid (R)-1-(3-amino-4-benzyloxy-phenyl)-2-[benzyl-(2,5,6-trimethyl-indan-2-yl)-amino]-ethyl ester

(R)-1-(4-Benzyloxy-3-nitro-phenyl)-2-[benzyl-(2,5,6-trimethyl-indan-2-yl)-amino]-ethanol

A mixture of 2-(4-methyl-3-nitro-phenyl)-oxirane and benzyl-(2,5,6-trimethyl-indan-2-yl)-amine is heated at 110°C for 48 hours. The material is used without further purification.

ES⁺ MS m/e 538 (MH⁺)

Acetic acid (R)-1-(4-benzyloxy-3-nitro-phenyl)-2-[benzyl-(2,5,6-trimethyl-indan-2-yl)-amino]-ethyl ester

To a solution of (R)-1-(4-Benzyloxy-3-nitro-phenyl)-2-[benzyl-(2,5,6-trimethyl-indan-2-yl)-amino]-ethanol in pyridine is added acetic anhydride and the mixture stirred for 18 hours.

The reaction mixture is quenched with water and after addition of ethyl acetate washed twice with aqueous KHSO₄ solution, twice with aqueous NaHCO₃ and once with brine.

The product is purified by chromatography (silica, ethyl acetate / hexane 1:4). ES⁺ MS m/e 579 (MH⁺)

Acetic acid (R)-1-(3-amino-4-benzyloxy-phenyl)-2-[benzyl-(2,5,6-trimethyl-indan-2-yl)-amino]-ethyl ester

Acetic acid (R)-1-(4-benzyloxy-3-nitro-phenyl)-2-[benzyl-(2,5,6-trimethyl-indan-2-yl)-amino]-ethyl ester in a mixture of THF and toluene is stirred under hydrogen in the presence of PtO₂ at room temperature for 15 hours. The mixture is filtered through celite and the filtrate is concentrated *in vacuo*. ES⁺ MS m/e 549 (MH⁺)

Intermediate 30 - 5,6-Diethyl-2-methyl-indan-2-ylamine

N-(5-Acetyl-2-methyl-indan-2-yl)-benzamide

Aluminium chloride (3.7g) is dissolved in nitromethane (12ml) under nitrogen followed by N-(2-methyl-indan-2-yl)-benzamide (3.0g) at 0°C. Acetyl chloride (0.85ml) is added dropwise over 30 minutes. After 4 hours at room temperature the mixture is quenched with ice and concentrated HCl, extracted with DCM. The organic layers are washed with dilute HCl and brine. Evaporation of the solvent yielded the desired product. ES⁺ MS m/e 294 (MH⁺)

N-(5-Ethyl-2-methyl-indan-2-yl)-benzamide

A solution of N-(5-acetyl-2-methyl-indan-2-yl)-benzamide (3.4g) in ethanol (200ml) and conc. HCl (2ml) is stirred under hydrogen in the presence of 10% Pd/C at room temperature for 48 hours. The mixture is filtered through celite and the filtrate is concentrated *in vacuo* to give the title compound.

¹H-NMR (CDCl₃) ppm: 1.20 (t, 3H), 1.60 (s, 3H), 2.55 (q, 2H), 3.05 (d, 2H), 3.35 (d, 2H), 6.35 (br.s, NH), 6.90-7.10 (m, 3H), 7.39 (d, 2H), 7.65 (s, 2H)

N-(5-Acetyl-6-ethyl-2-methyl-indan-2-yl)-benzamide is prepared from N-(5-ethyl-2-methyl-indan-2-yl)-benzamide (2.6g) following the procedure used to prepare N-(5-acetyl-2-methyl-indan-2-yl)-benzamide. The product is purified by chromatography (silica, hexane / ethyl acetate, 4:1) to give the title compound. ES⁺ MS m/e 322 (MH⁺)

N-(5,6-Diethyl-2-methyl-indan-2-yl)-benzamide is prepared from N-(5-acetyl-6-ethyl-2-methyl-indan-2-yl)-benzamide (1.1g) following the procedure used to prepare N-(5-ethyl-2-methyl-indan-2-yl)-benzamide. ES⁺ MS m/e 308 (MH⁺)

Benzyl-(5,6-diethyl-2-methyl-indan-2-yl)-amine is prepared from N-(5,6-diethyl-2-methyl-indan-2-yl)-benzamide by an analogous procedure to that used to prepare benzyl-(5,6-diethyl-indan-2-yl)-amine in Intermediate 18. ES⁺ MS m/e 294 (MH⁺)

5,6-Diethyl-2-methyl-indan-2-ylamine

A solution of benzyl-(5,6-diethyl-2-methyl-indan-2-yl)-amine (0.48g) in methanol (10ml) is stirred under an atmosphere of hydrogen in the presence of 10% Pd/C at room temperature for 18 hours. The mixture is filtered through celite and the filtrate is concentrated *in vacuo* to give the title compound. ES⁺ MS m/e 204 (MH⁺)

Example 1

(R)-8-Benzyloxy-5-[2-(4,7-dimethoxy-indan-2-ylamino)-1-hydroxy-ethyl]-1H-quinolin-2-one

(R)-8-Benzyloxy-5-oxiranylcarbostyryl (100mg, 0.34mmol), prepared from literature procedure (Beeley, Lee James; Dean, David Kenneth, PCT Int. Appl. WO 9525104)

and 4,7-dimethoxy-indan-2-ylamine (66mg, 0.34mmol), prepared from literature procedure (Sindelar, R. D.; Mott, J.; Barfknecht, C. F.; Arneric, S. P.; Flynn, J. R.; Long, J. P.; Bhatnagar, R. K. J. Med. Chem. (1982), 25(7), 858-64), are dissolved in toluene (1mL). The

reaction mixture was heated to 110°C and the solvent is allowed to evaporate. The residue is then stirred at 110°C for 4 hours. The reaction is shown to be complete by TLC. The product is purified by flash column chromatography (silica, dichloromethane / methanol 20:1).

TLC (silica, dichloromethane / methanol 25:1 $R_f = 0.10$).

ES+ MS m/e 487 (MH^+).

(R)-8-hydroxy-5-[2-(4,7-Dimethoxy-indan-2-ylamino)-1-hydroxy-ethyl]-1H-quinolin-2-one hydrochloride

(R)-8-Benzyloxy-5-[2-(4,7-dimethoxy-indan-2-ylamino)-1-hydroxy-ethyl]-1H-quinolin-2-one (37mg, 0.08mmol) is dissolved in methanol (10mL) and the compound is deprotected by adding a catalytic amount of 10% palladium on charcoal and placing the solution under an atmosphere of hydrogen. The reaction is shown to be complete by TLC after 4 hours. The catalyst is filtered off, 1M HCl/diethyl ether (1.1 equivalent) is added and the solvent is removed *in vacuo*.

TLC (silica, dichloromethane / methanol 10:1 $R_f = 0.15$).

ES+ MS m/e 397 (MH^+).

Other compounds of formula I are prepared from (R)-8-benzyloxy-5-oxiranylcarbostyryl ((R)-2-(4-benzyloxy-3-nitrophenyl)-oxirane (Intermediate 15) in Example 11) and the appropriate compound of formula XVII by procedures analogous to Example 1. These compounds, in which R^1 is OH, R^2 and R^3 are H, Ar is a group of formula III in which R^{29} , R^{30} and R^{31} are H (except in Example 11, where Ar is a group of formula XV in which R^{13} is H) and n is 1 (except in Example 9 where n is 2) are shown in the following table.

Example	R^4	R^5	R^6	R^7	ES+MS m/e (MH^+)
2	H	CH_3CH_2	CH_3CH_2	H	393
3	H	CH_3	CH_3	H	365
4	CH_3CH_2	H	H	CH_3CH_2	393
5	H	$-(CH_2)_4-$		H	391
6	H	$-O(CH_2)_2O-$		H	395
7	H	$CH_3(CH_2)_3$	$CH_3(CH_2)_3$	H	449
8	H	$CH_3(CH_2)_2$	$CH_3(CH_2)_2$	H	421
9	H	H	H	H	365
10	H	CH_3OCH_2	CH_3OCH_2	H	

11	H	CH ₃ CH ₂	CH ₃ CH ₂	H	341
----	---	---------------------------------	---------------------------------	---	-----

Example 10: ¹H-NMR (d₄-MeOH) ppm: 2.78 (2H, m), 2.9 (2H, m), 3.15 (2H, m), 3.28 (6H, s), 3.7 (1H, m), 4.55 (1H, br s), 5.15 (1H, m), 6.58 (1H, d), 6.9 (1H, d), 7.11 (2H, s), 7.15 (1H, s), 8.25 (1H, s).

Example 12

8-Hydroxy-5-[1-hydroxy-2-(indan-2-ylamino)-ethyl]-1H-quinolin-2-one

Intermediate 10 (18 mg, 0.054 mmol) is dissolved in methanol (2 mL) and cooled on ice. Sodium borohydride (6 mg, 0.12 mmol) is added over 2 hours. Concentrated HCl is then added until pH reaches 1, and the reaction mixture filtered. The filtrate is washed with methanol. The combined liquid phases are evaporated and redissolved in methanol twice. After removal of the methanol in vacuo, the residue is redissolved in water and the pH brought to 12 with 1N KOH. The solvent is removed in vacuo and the residue coevaporated twice with toluene. The residue is purified by flash chromatography (silica, CH₂Cl₂/methanol 8:2). ES+ MS *m/e* 337 (MH⁺).

Example 13

5-[2-(5,6-Dimethoxy-indan-2-ylamino)-1-hydroxy-ethyl]-8-hydroxy-1H-quinolin-2-one

This compound is prepared from Intermediate 11 by a procedure analogous to that of Example 12. ES+MS *m/e* 397 (MH⁺)

Example 14

5-[2-(5,6-Diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-8-hydroxy-3-methyl-1H-quinolin-2-one

This is prepared from Intermediate 13 (21mg) by the hydrogenation procedure for removal of the benzyl group used in Example 1.

¹H-NMR (d₄-CH₃OH) ppm 1.11 (t, 6H), 2.11 (s, 3H), 2.58 (q, 4H), 3.01-3.37 (m, 6H), 4.10-4.16 (m, 1H), 5.31-5.38 (m, 1H), 6.91 (d, 1H), 7.00 (s, 2H), 7.21 (d, 1H), 8.13 (s, 1H).

Example 15

5-[2-(5,6-Diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-8-methoxymethoxy-6-methyl-1H-quinolin-2-one

This is obtained from Intermediate 14 (20 mg) and 5,6-diethyl-indan-2-ylamine (72 mg) according to the procedure used for preparation of Intermediate 13.

¹H-NMR (CDCl₃) ppm: 1.14 (t, 6H), 2.30 (s, 3H), 2.51 (q, 4H), 2.64-3.16 (m, 6H), 3.41 (s, 3H), 3.60-3.68 (m, 1H), 5.18-5.25 (m, 3H), 6.50 (d, 1H), 7.89-7.94 (m, 3H), 8.68 (d, 2H), 9.15 (s, br, 1H).

5-[2-(5,6-Diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-8-hydroxy-6-methyl-1H-quinolin-2-one

3N Hydrochloric acid (1mL) is added to a solution of 5-[2-(5,6-Diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-8-methoxymethoxy-6-methyl-1H-quinolin-2-one (12 mg) in isopropanol (1mL) and tetrahydrofuran (1mL) at room temperature and the reaction mixture heated for 18 hours at 40°C. The solvent is removed *in vacuo*, and the product purified by preparative scale HPLC on a C8 column, eluting with a water/acetonitrile/trifluoroacetic acid gradient.

¹³C-NMR (d₄-CH₃OH) ppm: 15.97, 20.09, 26.34, 36.87, 51.75, 59.72, 67.33, 118.41, 119.12, 121.21, 125.45, 126.11, 128.60, 133.35, 137.52, 137.55, 142.32, 142.50, 145.69, 163.24.

Example 16**8-hydroxy-5-[2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-3,4-dihydro-1H-quinolin-2-one**

Hydrogenation of a methanol/ethanol solution of 8-hydroxy-5-[2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-1H-quinolin-2-one (Example 2) with a 10% palladium on carbon catalyst at 30°C for 48 hours under one atmosphere of hydrogen gives the title compound after filtration and evaporation. Further purification is achieved via preparative HPLC (column: Phenomenex Luna 10µm 150 mm x 50 mm, eluent: gradient from 10% to 95% acetonitrile in water containing 0.1% trifluoroacetic acid, UV detection at 254 nm).

¹³C-NMR (d₆-DMSO) ppm: 15.77, 21.42, 25.01, 30.37, 37.73, 37.83, 53.88, 58.68, 67.37, 113.28, 120.21, 122.08, 124.31, 124.34, 131.01, 138.46, 138.52, 139.58, 143.12, 169.44.

Example 17**(a) Acetic acid (R)-1-(4-benzyloxy-3-formylamino-phenyl)-2-[benzyl-(2,5,6-trimethyl-indan-2-yl)-amino]-ethyl ester**

To Intermediate 29 in toluene / THF is slowly added an aged mixture of formic acid and acetic anhydride and the reaction is stirred for 5 hours at room temperature. Ethyl acetate is added and washed with saturated NaHCO₃ solution. Purification by chromatography (silica, ethyl acetate / hexane 1:2) and trituration with ether gave off-white crystals. ES⁺ MS m/e 577 (MH⁺)

(b) N-(2-Benzyloxy-5-((R)-2-[benzyl-(2,5,6-trimethyl-indan-2-yl)-amino]-1-hydroxy-ethyl)-phenyl)-formamide

The product of Example 17(a) is suspended in ethanol and a catalytic amount of NaOCH₃ in methanol is added. After two hours at 70⁰C the solvent is removed and the residue purified by chromatography (silica, ethyl acetate / hexane 2:3) to give white crystals. ES⁺ MS m/e 535 (MH⁺)

(c) N-{2-Hydroxy-5-[(R)-1-hydroxy-2-(2,5,6-trimethyl-indan-2-ylamino)-ethyl]-phenyl}-formamide is prepared from the product of example 17(b) by a procedure analogous to that of Example 34(c). ES⁺ MS m/e 355 (MH⁺)Example 18**(a) 8-Benzylamino-5-[(R)-2-(5,6-diethyl-2-methyl-indan-2-ylamino)-1-hydroxy-ethyl]-1H-quinolin-2-one**

A mixture of 5,6-diethyl-2-methyl-indan-2-ylamine (0.28g) and 8-benzyloxy-5-oxiranyl-1H-quinolin-2-one (0.42g) in n-butanol (0.7ml) is placed in a Prolabo microwave oven for 75 minutes. at 100⁰C. The product is purified by chromatography (silica, DCM / ethanol, 5:1) to give the desired product. ES⁺ MS m/e 497 (MH⁺)

5-[(R)-2-(5,6-Diethyl-2-methyl-indan-2-ylamino)-1-hydroxy-ethyl]-8-hydroxy-1H-quinolin-2-one

A solution of the product of Example 18(a) (0.20g) in methanol (20ml) is stirred under an atmosphere of hydrogen in the presence of 10% Pd/C at room temperature for 2 hours. The

mixture is filtered through celite and the filtrate is concentrated *in vacuo*. Trituration with diethyl ether gave the desired product. ES⁺ MS m/e 407 (MH⁺)

Example 19

(a) (S)-8-Benzyloxy-5-[2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-1H-quinolin-2-one is prepared from Intermediate 16 (152mg) and Intermediate 1 (100mg) using a procedure analogous to that of Example 1(a). TLC (silica, dichloromethane / methanol 10:1 R_f = 0.25).

(b) (S)-5-[2-(4,7-Diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-1H-quinolin-2-one hydrochloride is prepared from the product of Example 19(a) by a procedure analogous to that of Example 1(b). TLC (silica, dichloromethane / methanol 10:1 R_f = 0.05).

Example 20

(a) 8-Benzyloxy-5-[(R)-1-hydroxy-2-(6,7,8,9-tetrahydro-5H-benzocyclohepten-7-ylamino)-ethyl]-1H-quinolin-2-one is prepared from (R)-8-benzyloxy-5-oxiranylcarbostyryl (203mg) and Intermediate 17 (110mg) by a procedure analogous to that of Example 1(a). TLC (silica, dichloromethane / methanol 10:1 R_f = 0.30).

(b) 5-[(R)-1-Hydroxy-2-(6,7,8,9-tetrahydro-5H-benzocyclohepten-7-ylamino)-ethyl]-1H-quinolin-2-one hydrochloride is prepared from the product of Example 20(a) by a procedure analogous to that of Example 1(b). TLC (silica, dichloromethane / methanol 10:1 R_f = 0.05).

Example 21

(a) (R)- 8-benzyloxy-5-[(S)-2-[benzyl-(5,6-diethyl-indan-2-yl)-amino]-1-hydroxy-ethyl]-1H-quinolin-2-one

A solution of (R)-8-benzyloxy-5-oxiranylcarbostyryl (5.00g) and 2-amino-5,6-diethylindan (3.87g) in n-butanol is heated for 4 hours at 110°C. After cooling to room temperature toluene (100ml) is added and the organic phase is washed with water (3 X 25ml), loaded onto a silica gel chromatography column and eluted with toluene followed by a mixture of toluene: ethanol: ethyl acetate: conc. ammonia (45:10:45:2) to give the title compound.

(b) (R)-5-[2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-1H-quinolin-2-one maleate (R)-8-benzyloxy-5-[2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-1H-quinolin-2-one (360mg) is dissolved in methanol (10mL) and the compound is deprotected by adding a catalytic amount of 10% palladium on charcoal and placing the solution under an atmosphere of hydrogen. The reaction is shown to be complete by TLC after 4 hours. The catalyst is filtered off and the solvent is removed *in vacuo*. The product is taken up into isopropanol and a solution of maleic acid in isopropanol added. The title compound is obtained after recrystallisation from ethanol. TLC (silica, dichloromethane / methanol 10:1 $R_f = 0.05$). ES+ MS *m/e* 393 (MH^+).

Example 22

(a) N-(5-[(R)-2-[Benzyl-(5,6-diethyl-indan-2-yl)-amino]-1-hydroxy-ethyl]-2-benzyloxy-phenyl)-formamide is prepared from Intermediate 19 (1.00g), formic acid (155mg) and acetic anhydride (226mg) using a procedure analogous to that of Example 21(a). TLC (silica, n-hexane / ethyl acetate 2:1 $R_f = 0.20$).

(b) N-{5-[(R)-2-(5,6-Diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-2-hydroxy-phenyl}-formamide is prepared from the product of Example 22(a) by a procedure analogous to that of Example 1(b). TLC (silica, n-hexane / ethyl acetate 2:1 $R_f = 0.05$).

Example 23

(a) (R)-2-[Benzyl-(5,6-diethyl-indan-2-yl)-amino]-1-(4-benzyloxy-3-dimethylamino-phenyl)-ethanol

Intermediate 19 (0.37g) is dissolved in CH_3OH (50mL) and formaldehyde, 37% in water (5mL), dissolved in water (10mL), is added. A catalytic amount of PtO_2 is added and the solution is stirred under an atmosphere of H_2 . The reaction is shown to be complete by TLC after 24 hours. The catalyst is filtered off, the solvent is removed *in vacuo* and the residue is partitioned between ethyl acetate (100mL) and water (100mL). The organic layer is dried over $MgSO_4$, filtered and the solvent is removed *in vacuo*. The product is purified by flash column chromatography (silica, n-hexane / ethyl acetate 4:1). TLC (silica, n-hexane / ethyl acetate 2:1 $R_f = 0.65$).

(b) 4-[(R)-2-(5,6-Diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-2-dimethylamino-phenol hydrochloride is prepared from the product of Example 23(a) by a procedure analogous to that of Example 1(b).

¹H NMR [DMSO , 400MHz] δ 1.10(6H,t), 2.55 (4H, q), 3.05(2H,m), 3.10 (6H, s), 3.20 (4H, m), 4.00 (1H, m), 4.95 (1H, m), 7.00 (2H, s), 7.15 (1H, d), 7.35 (1H, d), 7.80 (1H, s), 9.20 (1H, broad), 9.75 (1H, broad), 11.40 (1H, broad).

Example 24

(a) (R)-2-[Benzyl-(5,6-diethyl-indan-2-yl)-amino]-1-(4-benzyloxy-3-methylamino-phenyl)-ethanol

The product of Example 22 (260mg) is dissolved in dioxan (20mL). Sodium borohydride (90mg) is added followed by the dropwise addition of acetic anhydride (142mg). The reaction mixture is stirred at 90°C. The reaction is shown to be complete by TLC after 4 hours. The solvent is removed *in vacuo* and the residue is partitioned between ethyl acetate (100mL) and water (100mL). The organic layer is dried over MgSO₄, filtered and the solvent is removed *in vacuo*. The product is purified by flash column chromatography (silica, n-hexane / ethyl acetate 4:1). TLC (silica, n-hexane / ethyl acetate 2:1 R_f = 0.65).

(b) 4-[(R)-2-(5,6-Diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-2-methylamino-phenol hydrochloride is prepared from the product of Example 24(a) by a procedure analogous to that of Example 1(b).

¹H NMR [DMSO , 400MHz] δ 1.10(6H,t), 2.55 (4H, q), 2.85 (3H, s), 3.10 (6H, m), 4.00 (1H, m), 4.90 (1H, m), 7.00 (3H, m), 7.15 (1H, m), 7.40 (1H, m), 9.10 (1H, broad), 9.60 (1H, broad), 10.80 (1H, broad).

Example 25

(a) N-(5-[[Benzyl-(5,6-diethyl-indan-2-yl)-amino]-acetyl]-2-benzyloxy-phenyl)-methanesulfonamide

Intermediate 20 (240mg) is dissolved in dichloromethane (10mL). Triethylamine (56mg) is added followed by methanesulfonyl chloride (58mg). The reaction mixture is stirred at room temperature. The reaction is shown to be complete by TLC after 24 hours. The solvent is removed *in vacuo* and the product is purified by flash column chromatography (silica, n-hexane / ethyl acetate 4:1). TLC (silica, n-hexane / ethyl acetate 2:1 R_f = 0.40).

(b) **N-(5-{2-[Benzyl-(5,6-diethyl-indan-2-yl)-amino]-1-hydroxy-ethyl}-2-benzyloxy-phenyl)-methanesulfonamide**

The product of Example 25(a) (120mg) is dissolved in ethanol (10mL). Sodium borohydride (9mg) is added and the reaction mixture is stirred at room temperature. The reaction is shown to be complete by TLC after 3 hours. The reaction mixture is quenched with 2M HCl (1mL), the solvent is removed *in vacuo* and the residue is partitioned between ethyl acetate (50mL) and saturated NaHCO₃ (50mL). The organic layer is dried over MgSO₄, filtered and the solvent is removed *in vacuo*. The product is not purified further. TLC (silica, n-hexane / ethyl acetate 2:1 R_f = 0.45).

(c) **N-{5-[2-(5,6-Diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-2-hydroxy-phenyl}-methanesulfonamide hydrochloride** is prepared from the product of Example 25(b) by a procedure analogous to that of Example 1(b).

¹H NMR [CDCl₃, 400MHz] δ 1.15(6H,t), 2.55 (4H, q) 2.95 (3H,s), 3.10 (6H,m), 4.00 (1H, m), 4.85 (1H, m), 6.10 (1H, broad), 6.90 (2H, d), 7.00 (2H, s), 7.10 (1H, d of d), 7.25 (1H, d), 8.75 (1H, s), 8.95 (1H, broad), 9.25 (1H, broad), 10.00 (1H, s).

Example 26

(a) **(R)-8-Benzyloxy-5-[(S)-2-[benzyl-(4,5,6,7-tetramethyl-indan-2-yl)-amino]-1-hydroxy-ethyl]-1H-quinolin-2-one**

(R)-8-Benzyloxy-5-oxiranylcabostyryl (204 mg) and Intermediate 21 (194 mg) are dissolved in n-butanol (0.5ml) under nitrogen. The reaction mixture is heated at 110°C for 22 hours. On cooling the solvent is removed in vacuo. The product is purified by flash column chromatography (silica, ethyl acetate/heaxane 50:50). ES⁺ MS m/e 573 (MH⁺)

(b) **(R)-8-Hydroxy-5-[(S)-1-hydroxy-2-(4,5,6,7-tetramethyl-indan-2-ylamino)-ethyl]-1H-quinolin-2-one** is prepared from the product of Example 26(a) by a procedure analogous to that of Example 1(b).

¹H NMR (CD₃OD) ppm: 8.55 (1H, d); 7.5 (1H, d); 7.25 (1H, d); 6.9 (1H, d); 5.6 (1H, m); 4.3 (1H, m); 3.7 (2H, q); 3.6 (2H, dd); 3.3 (2H, dd); 2.4 (12H, s)

Example 27

(a) **8-Benzyloxy-5-[(R)-1-hydroxy-2-(2-methyl-indan-2-ylamino)-ethyl]-1H-quinolin-2-one.**

A mixture of 8-benzyloxy-5-(R)-oxiranyl-1H-quinolin-2-one (500 mg) and 2-methyl-indan-2-ylamine (276 mg) in n-butanol (1 mL) is subjected to microwave irradiation, using a Prolabo Synthwave 402 instrument, for 90 minutes at 110°C. The residue is absorbed on silica and the product is purified by flash chromatography (silica, chloroform/ethanol 4:1).

¹H-NMR (CDCl₃) ppm: 1.30 (s, 3H), 2.65 (s, 1H), 2.95 (dd, 2H), 3.07 (m, 3H), 5.15 (m, 1H), 5.18 (s, 2H), 6.66 (d, 1H), 7.17 (m, 4H), 7.26 (d, 1H), 7.45 (m, 5H), 8.07 (d, 1H), 8.8-9.5 (br.d, 1H)

(b) **8-Hydroxy-5-[(R)-1-hydroxy-2-(2-methyl-indan-2-ylamino)-ethyl]-1H-quinolin-2-one**

The product of Example 27(a) (100 mg, 0.22 mmol) is dissolved in methanol (20 mL) and is deprotected by adding a catalytic amount of 10% palladium on charcoal and stirring under an atmosphere of hydrogen for 1 hour. The catalyst is removed and the solvent is evaporated to give a yellow solid.

¹H-NMR (d₄-CH₃OH) ppm: 1.20 (s, 3H), 2.75 (m, 4H), 2.95 (d, 2H), 5.03 (m, 1H), 6.60 (d, 1H), 6.82 (d, 1H), 7.0 (m, 4H), 7.08 (d, 1H), 8.20 (d, 1H).

Example 28

5-[2-(5,6-Diethyl-indan-2-ylamino)-ethyl]-8-hydroxy-1H-quinolin-2-one

This compound is prepared from the product of Example 2 according to the procedure of Temple et al, J. Med. Chem., 19, 626-633 (1976).

¹H-NMR (d₄-CH₃OH) ppm: 1.08 (t, 3H), 2.55 (q, 4H), 2.96 (dd, 2H), 3.18 (m, 4H), 3.28 (dd, 2H), 3.99 (m, 1H), 6.60 (d, 1H), 6.90 (d, 1H), 6.97 (d, 1H), 6.00 (s, 2H), 8.07 (d, 1H).

Example 29

(a) **8-Benzyloxy-5-[(R)-1-hydroxy-2-(2-methyl-2,3,5,6,7,8-hexahydro-1H-cyclopenta[b]naphthalen-2-ylamino)-ethyl]-1H-quinolin-2-one** is prepared from 8-benzyloxy-5-(R)-oxiranyl-1H-quinolin-2-one (220 mg) and Intermediate 24 (150 mg) by procedures analogous to those of Example 27(a).

¹H-NMR (CDCl₃) ppm: 1.37 (s, 1H), 1.78 (m, 4H), 2.1 (br.s, 2H), 2.72 (m, 5H), 2.80 (dd, 2H), 2.95 (m, 3H), 5.08 (m, 1H), 5.17 (s, 2H), 6.65 (d, 1H), 6.88 (s, 2H), 7.02 (d, 2H), 7.26 (d, 1H), 7.4 (m, 5H), 8.05 (d, 1H).

(b) 8-Hydroxy-5-[(R)-1-hydroxy-2-(2-methyl-2,3,5,6,7,8-hexahydro-1H-cyclopenta[b]naphthalen-2-ylamino)-ethyl]-1H-quinolin-2-one is prepared by hydrogenation of the product of Example 29(a) using a procedure analogous to that of Example 27(b). The product is purified by HPLC (H₂O, CH₃CN, CF₃COOH, gradient elution).

¹H-NMR (d₄-CH₃OH) ppm (TFA salt): 1.65 (s, 3H), 1.85 (m, 4H), 2.85 (m, 4H), 3.15 (m, 2H), 3.4 (m, 4H), 5.48 (t, 1H), 6.83 (d, 1H), 7.03 (s, 2H), 7.15 (d, 1H), 7.45 (d, 1H), 8.40 (d, 1H).

Example 30

(a) 5-[(S)-2-[Benzyl-(2,3,5,6,7,8-hexahydro-1H-cyclopenta[b]naphthalen-2-yl)-amino]-1-hydroxy-ethyl]-8-benzyloxy-1H-quinolin-2-one

A mixture of Intermediate 16 (150mg) and benzyl-(2,3,5,6,7,8-hexahydro-1H-cyclopenta[b]naphthalen-2-yl)-amine (142 mg) in toluene (1 mL) is heated at 80°C for 36 hours. The residue is purified by chromatography (silica, CHCl₃ / EtOH, 20:1) to give a yellow foam.

¹H-NMR (CDCl₃) ppm: 1.77 (m, 4H), 2.72 (m, 6H), 3.01 (m, 4H), 3.70 (d, 1H), 3.88 (d, 1H), 4.82 (m, 1H), 5.15 (s, 2H), 6.50 (d, 1H), 6.8 - 8 (m, 13H), 9.05 (br.s, 1H)

(b) 5-[(S)-2-(2,3,5,6,7,8-Hexahydro-1H-cyclopenta[b]naphthalen-2-ylamino)-1-hydroxy-ethyl]-8-hydroxy-1H-quinolin-2-one

A solution of the product of Example 30(a) (150 mg) in methanol (20 mL) is stirred under an atmosphere of hydrogen in the presence of 10% Pd/C (20 mg) at room temperature for 5 hours. The reaction is filtered and the product is purified by chromatography (silica, CHCl₃ / EtOH, 20:1) followed by crystallisation (CH₃OH).

¹H-NMR (d₄-CH₃OH) ppm: 1.65 (m, 4H), 2.57 (m, 4H), 2.86 (dd, 2H), 3.1 (m, 4H), 3.82 (m, 1H), 5.25 (m, 1H), 6.55 (d, 1H), 6.78 (s, 2H), 6.91 (d, 1H), 7.19 (d, 1H), 8.27 (d, 1H).

Example 31

(a) Acetic acid (R)-2-[benzyl-(2-methyl-indan-2-yl)-amino]-1-(4-benzyloxy-3-methanesulfonylamino-phenyl)-ethyl ester is prepared from Intermediate 27 (476mg), triethylamine (231mg) and methanesulfonyl chloride (210mg) by a procedure analogous to that of Example 25(b). TLC (silica, n-hexane / ethyl acetate 2:1 R_f = 0.45).

(b) N-(5-((R)-2-[Benzyl-(2-methyl-indan-2-yl)-amino]-1-hydroxy-ethyl)-2-benzyloxy-phenyl)-methanesulfonamide

The product of Example 31(a) (200mg) is dissolved in CH₃OH (8mL). K₂CO₃ (138mg) is added followed by the dropwise addition of water (2mL). The reaction mixture is stirred at room temperature. The reaction is shown to be complete by TLC after 24 hours. Ethyl acetate (100mL) is added and the solution is washed with water (50mL) and brine (50mL). The organic layer is dried over MgSO₄, filtered and the solvent is removed *in vacuo*. The product is purified by flash column chromatography (silica, n-hexane / ethyl acetate 3:1). TLC (silica, n-hexane / ethyl acetate 2:1 R_f = 0.35).

(c) N-{2-Hydroxy-5-[(R)-1-hydroxy-2-(2-methyl-indan-2-ylamino)-ethyl]-phenyl}-methanesulfonamide is prepared from the product of Example 31(b) by a procedure analogous to that of Example 1(b). TLC (silica, dichloromethane / methanol 10:1 R_f = 0.10).

Example 32

(a) Acetic acid (R)-2-[benzyl-(2-methyl-indan-2-yl)-amino]-1-(4-benzyloxy-3-ethanesulfonylamino-phenyl)-ethyl ester is prepared from Intermediate 27, triethylamine (242mg) and ethanesulfonyl chloride (247mg) by a procedure analogous to that of Example 25(b). TLC (silica, n-hexane / ethyl acetate 2:1 R_f = 0.50).

(b) Ethanesulfonic acid (5-((R)-2-[benzyl-(2-methyl-indan-2-yl)-amino]-1-hydroxy-ethyl)-2-benzyloxy-phenyl)-amide is prepared from the product of Example 32(a) by a procedure analogous to that in Example 31(b). TLC (silica, n-hexane / ethyl acetate 2:1 R_f = 0.40).

(c) Ethanesulfonic acid {2-hydroxy-5-[(R)-1-hydroxy-2-(2-methyl-indan-2-ylamino)-ethyl]-phenyl}-amide is prepared from the product of Example 32(b) by a procedure analogous to that of Example 1(b). TLC (silica, dichloromethane / methanol 10:1 R_f = 0.10).

Example 33

(a) Acetic acid (R)-2-[benzyl-(2-methyl-indan-2-yl)-amino]-1-[4-benzyloxy-3-(propane-1-sulfonylamino)-phenyl]-ethyl ester is prepared from Intermediate 27 (525mg), triethylamine

(255mg) and 1-propanesulfonyl chloride (288mg) by a procedure analogous to that of Example 25(a). TLC (silica, n-hexane / ethyl acetate 4:1 R_f = 0.25).

(b) Propane-1-sulfonic acid (5-[(R)-2-[benzyl-(2-methyl-indan-2-yl)-amino]-1-hydroxy-ethyl]-2-benzyloxy-phenyl)-amide is prepared from the product of Example 33(a) by a procedure analogous to of Example 31(b). TLC (silica, n-hexane / ethyl acetate 4:1 R_f = 0.15).

(c) Propane-1-sulfonic acid [2-hydroxy-5-[(R)-1-hydroxy-2-(2-methyl-indan-2-ylamino)-ethyl]-phenyl]-amide is prepared from the product of Example 33(b) by a procedure analogous to that of Example 1(b). TLC (silica, dichloromethane / methanol 10:1 R_f = 0.05).

Example 34

(a) N-[2-Benzyloxy-5-[(2-ethyl-indan-2-ylamino)-acetyl]-phenyl]-methanesulfonamide

A mixture of 2-ethyl-indan-2-ylamine and N-(2-benzyloxy-5-bromoacetyl-phenyl)-methanesulfonamide is stirred in acetonitrile at room temperature for 20 hours. The product is isolated by filtration. ES⁺ MS m/e 479 (MH⁺)

(b) N-[2-Benzyloxy-5-[2-(2-ethyl-indan-2-ylamino)-1-hydroxy-ethyl]-phenyl]-methanesulfonamide

The product of Example 34(a) is suspended in a mixture of ethanol and dichloromethane. Sodium borohydride is added at 0°C and the mixture stirred at room temperature for 3 hours, then filtered and chromatographed (silica, ethylacetate / ethanol 4:1) to give a white foam. ES⁺ MS m/e 480 (MH⁺)

(c) N-[5-[2-(2-Ethyl-indan-2-ylamino)-1-hydroxy-ethyl]-2-hydroxy-phenyl]-methanesulfonamide

The product of Example 34(b) (0.29 g) in methanol (20 mL) is stirred under hydrogen in the presence of 10% Pd/C at room temperature for 18 hours. The mixture is filtered through celite and the filtrate is concentrated *in vacuo*, then chromatographed (silica, ethyl acetate / ethanol 2:1). After trituration with ether / ethyl acetate off-white crystals (100 mg) are obtained.

¹H-NMR (d₄-CH₃OH) ppm: 0.85 (t, 3H), 1.65 (m, 2H), 2.75 (m, 2H), 2.85 (s, 3H), 2.95 (m, 4H), 6.80 (d, 1H), 7.05 (m, 5H), 7.30 (s, 1H).

ES⁺ MS m/e 491 (MH⁺)

Example 35

(a) **Acetic acid (R)-1-(4-benzyloxy-3-methanesulfonylamino-phenyl)-2-[benzyl-(2,5,6-trimethyl-indan-2-yl)-amino]-ethyl ester**

To a solution of Intermediate 29 in dichloromethane and triethylamine at room temperature is added methanesulfonyl chloride and the mixture stirred for 18 hours. It is then washed with 0.2 N HCl, saturated NaHCO₃ solution and brine. The product is purified by chromatography (silica, ethyl acetate / hexane 1:4). ES⁺ MS m/e 625 (M⁺)

(b) **N-(2-Benzyloxy-5-[(R)-2-[benzyl-(2,5,6-trimethyl-indan-2-yl)-amino]-1-hydroxy-ethyl]-phenyl)-methanesulfonamide**


The product of Example 35(a) is stirred in methanol / water with K₂CO₃ for 3 days then solvents removed *in vacuo*. The product is purified by chromatography (silica, ethyl acetate / hexane 1:2).

¹H-NMR (CDCl₃) ppm: 1.21 (s, 3H), 2.22 (s, 6H), 2.63-2.82 (m, 4H), 2.84 (s, 3H), 3.20 (br.d, 2H), 3.61 (d, 1H), 3.64 (br.s., 1H), 3.83 (m, 1H), 4.08 (d, 1H), 5.09 (s, 2H), 6.75 (br.s, NH), 6.90-7.05 (m, 4H), 7.25- 7.45 (11H).

(c) **N-{2-Hydroxy-5-[(R)-1-hydroxy-2-(2,5,6-trimethyl-indan-2-ylamino)-ethyl]-phenyl}-methanesulfonamide** is prepared from the product of Example 35(b) by a procedure analogous to that of Example 34(c). ES⁺ MS m/e 405 (MH⁺)

Claims


1. A compound of formula



I

in free or salt or solvate form, where

Ar is a group of formula



II

R¹ is hydrogen, hydroxy, or alkoxy,

R² and R³ are each independently hydrogen or alkyl,

R⁴, R⁵, R⁶ and R⁷ are each independently hydrogen, halogen, cyano, hydroxy, alkoxy, aryl, alkyl, alkyl substituted by one or more halogen atoms or one or more hydroxy or alkoxy groups, alkyl interrupted by one or more hetero atoms, alkenyl, trialkylsilyl, carboxy, alkoxycarbonyl, or -CONR¹¹R¹², where R¹¹ and R¹² are each independently hydrogen or alkyl, or R⁴ and R⁵, R⁵ and R⁶, or R⁶ and R⁷ together with the carbon atoms to which they are attached denote a carbocyclic or heterocyclic ring,

R⁸ is halogen, -OR¹³, -CH₂OR¹³ or -NHR¹³ where R¹³ is hydrogen, alkyl, alkyl interrupted by one or more hetero atoms, -COR¹⁴, where R¹⁴ is hydrogen, -N(R¹⁵)R¹⁶, alkyl or alkyl interrupted by one or more hetero atoms, or aryl and R¹⁵ and R¹⁶ are each independently hydrogen, alkyl or alkyl interrupted by one or more hetero atoms, or R¹³ is -C(=NH)R¹⁷, -SOR¹⁷ or -SO₂R¹⁷ where R¹⁷ is alkyl or alkyl interrupted by one or more hetero atoms, and R⁹ is hydrogen, or R⁸ is -NHR¹⁸ where -NHR¹⁸ and R⁹, together with the carbon atoms to which they are attached, denote a 5- or 6- membered heterocycle,

R^{10} is $-OR^{19}$ or $-NHR^{19}$ where R^{19} is hydrogen, alkyl, alkyl interrupted by one or more hetero atoms, or $-COR^{20}$, where R^{20} is $-N(R^{21})R^{22}$, alkyl or alkyl interrupted by one or more hetero atoms, or aryl, and R^{21} and R^{22} are each independently hydrogen, alkyl or alkyl interrupted by one or more hetero atoms,

X is halogen or halomethyl or alkyl,

Y is carbon or nitrogen,

n is 1 or 2,

p is zero when Y is nitrogen or 1 when Y is carbon,

q and r are each zero or 1, the sum of q+r is 1 or 2; and

the carbon atom marked with an asterisk* has the R or S configuration, or a mixture thereof, when R^1 is hydroxy or alkoxy.

2. A compound according to claim 1, in which Ar is a group of formula II in which

Y is carbon,

R^8 is $-NHR^{18}$ and $-NHR^{18}$ and R^9 together denote

a group of formula $-NH-CO-R^{23}$ - where R^{23} is an alkylene, alkenylene or alkyleneoxy group,

a group of formula $-NH-SO_2-R^{24}$ where R^{24} is an alkyleneoxy group,

a group of formula $-NH-R^{25} (COOR^{26})-$ where R^{25} is an alkylene or alkenylene group and R^{26} is alkyl, or

a group of formula $-NH-CO-NH-$ or $-NH-CO-S-$,


R^{10} is $-OR^{19}$, where R^{19} is as defined in claim 1,

X is alkyl,





p is 1, q is 1 and r is zero or 1.

3. A compound according to claim 2, in which Ar is a group of formula III, IV, V, VI or VII

:




in which R^{29} , R^{30} and R^{31} are each independently hydrogen or C_1 - C_4 -alkyl



in which Z is -O-, -NH- or -S-.


4. A compound according to claim 1, in which Ar is a group of formula




where R^{29} , R^{30} and R^{31} are each independently hydrogen or C_1 - C_4 -alkyl.

5. A compound according to claim 1, in which Ar is a group of formula II in which Y is carbon, R^8 is $-CH_2OR^{13}$ where R^{13} is hydrogen, C_1 - C_4 -alkyl, or C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, R^9 is hydrogen, R^{10} is $-OR^{19}$ where R^{19} is hydrogen, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl or R^{10} is $-NHR^{19}$ where R^{19} is hydrogen, C_1 - C_4 -alkyl or $-COR^{20}$ where R^{20} is C_1 - C_4 -alkyl, C_6 - C_{10} -aryl or $-N(R^{21})R^{22}$ where R^{21} and R^{22} are each independently hydrogen or C_1 - C_4 -alkyl, p and q are each 1 and r is zero; or a group of formula II in which Y is nitrogen, R^8 is $-CH_2OR^{13}$ where R^{13} is hydrogen, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, R^{10} is $-OR^{19}$ where R^{19} is hydrogen, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, p and r are zero and q is 1.


6. A compound according to claim 5, in which Ar is a group of formula XII, XIII or XIV



XII




XIII



XIV

7. A compound according to claim 1, in which Ar is a group of formula II in which Y is carbon, R^8 is $-NHR^{13}$ where R^{13} is hydrogen, C_1 - C_{10} alkyl, C_1 - C_{10} alkyl interrupted by 1 to 3 hetero atoms, $-COR^{14}$ where R^{14} is hydrogen, C_1 - C_{10} -alkyl or C_1 - C_{10} -alkyl interrupted by 1 to 3 hetero atoms, or R^{13} is $-C(=NH)R^{17}$, $-SOR^{17}$ or $-SO_2R^{17}$ where R^{17} is C_1 - C_{10} -alkyl or C_1 - C_{10} -alkyl interrupted by 1 to 3 hetero atoms, R^9 is hydrogen, R^{10} is $-OR^{18}$ where R^{18} is hydrogen, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy- C_1 - C_4 alkyl, p and q are each 1 and r is zero.

8. A compound according to claim 7, in which Ar is a group of formula XV




XV

9. A compound according to any one of the preceding claims, in which R^4 , R^5 , R^6 and R^7 are each hydrogen or are such that the benzene ring to which they are attached is symmetrically substituted.

10. A compound according to claim 1, in which Ar is a group of formula III, IV, V, XII or XV, R^1 is hydroxy, R^2 and R^3 are hydrogen, and R^4 and R^7 are identical and are each hydrogen, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy, and either R^5 and R^6 are identical and are each hydrogen, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy or C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, or R^5 and R^6 together denote $-(CH_2)_4-$ or $-O(CH_2)_2O-$, in free or salt or solvate form.

11. A compound according to claim 10, in which the carbon atom in formula I marked with an asterisk * has the R configuration.


12. A compound of formula



I

in free or salt or solvate form,

(A) wherein Ar is a group of formula




III

in which R^{29} , R^{30} and R^{31} are each H, R^1 is OH, R^2 and R^3 are each H and

- (i) n is 1, and R^4 and R^7 are each $\text{CH}_3\text{O}-$ and R^5 and R^6 are each H; or
- (ii) n is 1, and R^4 and R^7 are each H and R^5 and R^6 are each CH_3CH_2- ; or
- (iii) n is 1, and R^4 and R^7 are each H and R^5 and R^6 are each CH_3- ; or
- (iv) n is 1, and R^4 and R^7 are each CH_3CH_2- and R^5 and R^6 are each H; or
- (v) n is 1, and R^4 and R^7 are each H and R^5 and R^6 together denote $-(\text{CH}_2)_4-$; or
- (vi) n is 1, and R^4 and R^7 are each H and R^5 and R^6 together denote $-\text{O}(\text{CH}_2)_2\text{O}-$; or
- (vii) n is 1, and R^4 and R^7 are each H and R^5 and R^6 are each $\text{CH}_3(\text{CH}_2)_3-$; or
- (viii) n is 1, and R^4 and R^7 are each H and R^5 and R^6 are each $\text{CH}_3(\text{CH}_2)_2-$; or
- (ix) n is 2, R^4 , R^5 , R^6 and R^7 are each H; or
- (x) n is 1, and R^4 and R^7 are each H and R^5 and R^6 are each CH_3OCH_2- ; or

(B) wherein Ar is a group of formula



XV

in which R^{13} is H, R^1 is OH, R^2 and R^3 are each H, R^4 and R^7 are each H and R^5 and R^6 are each H and n is 1; or

(C) which is a compound selected from 8-hydroxy-5-[1-hydroxy-2-(indan-2-ylamino)-ethyl]-1H-quinolin-2-one, 5-[2-(5,6-dimethoxy-indan-2-ylamino)-1-hydroxy-ethyl]-8-hydroxy-1H-quinolin-2-one, 5-[2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-8-hydroxy-3-methyl-1H-quinolin-2-one, 5-[2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-8-methoxymethoxy-6-methyl-1H-quinolin-2-one, 5-[2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-8-hydroxy-6-methyl-1H-quinolin-2-one, 8-hydroxy-5-[2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-3,4-dihydro-1H-quinolin-2-one, N-{2-hydroxy-5-[(R)-1-hydroxy-2-(2,5,6-trimethyl-indan-2-ylamino)-ethyl]-phenyl}-formamide, 5-[(R)-2-(5,6-diethyl-2-methyl-indan-2-ylamino)-1-hydroxy-ethyl]-8-hydroxy-1H-quinolin-2-one, (S)-5-[2-(4,7-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-1H-quinolin-2-one hydrochloride, 5-[(R)-1-hydroxy-2-(6,7,8,9-tetrahydro-5H-benzocyclohepten-7-ylamino)-ethyl]-1H-quinolin-2-one hydrochloride, (R)-5-[2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-1H-quinolin-2-one hydrochloride, N-{5-[(R)-2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-2-hydroxy-phenyl}-formamide, 4-[(R)-2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-2-dimethylamino-phenol hydrochloride, 4-[(R)-2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-2-methylamino-phenol hydrochloride, N-{5-[2-(5,6-diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-2-hydroxy-phenyl}-methanesulfonamide hydrochloride, (R)-8-hydroxy-5-[(S)-1-hydroxy-2-(4,5,6,7-tetramethyl-indan-2-ylamino)-ethyl]-1H-quinolin-2-one, 8-hydroxy-5-[(R)-1-hydroxy-2-(2-methyl-indan-2-ylamino)-ethyl]-1H-quinolin-2-one, 5-[2-(5,6-diethyl-indan-2-ylamino)-ethyl]-8-hydroxy-1H-quinolin-2-one, 8-hydroxy-5-[(R)-1-hydroxy-2-(2-methyl-2,3,5,6,7,8-hexahydro-1H-cyclopenta[b]naphthalen-2-ylamino)-ethyl]-1H-quinolin-2-one, 5-[(S)-2-(2,3,5,6,7,8-hexahydro-1H-cyclopenta[b]naphthalen-2-ylamino)-1-hydroxy-ethyl]-8-hydroxy-1H-quinolin-2-one, N-{2-hydroxy-5-[(R)-1-hydroxy-2-(2-methyl-indan-2-ylamino)-ethyl]-phenyl}-methanesulfonamide, ethanesulfonic acid {2-hydroxy-5-[(R)-1-hydroxy-2-(2-methyl-indan-2-ylamino)-ethyl]-phenyl}-amide, propane-1-sulfonic acid {2-hydroxy-5-[(R)-1-hydroxy-2-(2-methyl-indan-2-ylamino)-ethyl]-phenyl}-amide, N-{5-[2-(2-ethyl-indan-2-ylamino)-1-hydroxy-ethyl]-2-hydroxy-phenyl}-methanesulfonamide, or N-{2-hydroxy-5-[(R)-1-hydroxy-2-(2,5,6-trimethyl-indan-2-ylamino)-ethyl]-phenyl}-methanesulfonamide.

13. A pharmaceutical composition comprising a compound according to any one of the preceding claims, optionally together with a pharmaceutically acceptable carrier.


14. Use of a compound according to any one of claims 1 to 12 for the preparation of a medicament for the treatment of a condition which is prevented or alleviated by activation of the β 2-adrenoreceptor.

15. Use of a compound according to any one of claims 1 to 12 for the preparation of a medicament for the treatment of an obstructive or inflammatory airways disease.


16. A process for the preparation of a compound of formula I in free or salt or solvate form comprising:

(a) for the preparation of a compound where R^1 is hydroxy, either

(i) reacting a compound of formula




with a compound of formula



where Ar^1 is Ar as defined in claim 1 or a protected form thereof, R^2 , R^3 , R^4 , R^5 , R^6 , R^7 and n are as defined in claim 1 and R^{32} is hydrogen or an amine-protective group, or

(ii) reducing a compound of formula



where Ar¹ is Ar as defined in claim 1 or a protected form thereof, R², R³, R⁴, R⁵, R⁶ and R⁷ are as defined in claim 1, to convert the indicated keto group into -CH(OH)-; or

(b) for the preparation of a compound where R¹ is hydrogen, reducing a corresponding compound of formula I where R¹ is hydroxy; or

(c) for the preparation of a compound of formula I where R¹ is alkoxy, either (i) O-alkylating a corresponding compound of formula I where R¹ is hydroxy or (ii) reacting a corresponding compound having a leaving moiety instead of R¹ with an alcohol of formula R¹H where R¹ is alkoxy;

and, optionally, converting a resultant compound of formula I in protected form into a corresponding compound in unprotected form;

and recovering the resultant compound of formula I in free or salt or solvate form.

INTERNATIONAL SEARCH REPORT

Int. National Application No.
PCT/EP 00/05058

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D215/26 A61K31/395 A61P43/00 C07D405/12 C07C233/43
C07D215/22 C07C215/80 C07C311/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 882 704 A (KISSEI PHARMACEUTICAL CO., LTD) 9 December 1998 (1998-12-09) page 2, line 5 - line 24 page 3, line 25 - line 31; claim 1 ---	1,13-15
Y	WO 99 09001 A (KISSEI PHARMACEUTICAL CO., LTD) 25 February 1999 (1999-02-25) * abstract *	1,13-15
Y	WO 93 18007 A (TOKYO TANABE COMPANY LTD.) 16 September 1993 (1993-09-16) * abstract *	1,13-15
A	EP 0 894 787 A (KISSEI PHARMACEUTICAL CO., LTD) 3 February 1999 (1999-02-03) page 4 --- -/--	1,13-15



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

28 September 2000

Date of mailing of the international search report

18/10/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

Int. Application No.

PCT/EP 00/05058

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 93 15041 A (FUJISAWA PHARMACEUTICAL CO.,LTD.) 5 August 1993 (1993-08-05) page 35, line 30 -page 36, line 12; claim 1 -----	1,13
A	WO 92 18461 A (FUJISAWA PHARMACEUTICAL CO.,LTD.) 29 October 1992 (1992-10-29) page 1 -page 2; claim 1 -----	1,13

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. J. Application No

PCT/EP 00/05058

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 882704	A	09-12-1998	AU 2001497 A	02-09-1997
			BR 9707566 A	27-07-1999
			NO 983777 A	19-10-1998
			CA 2245490 A	21-08-1997
			CN 1216526 A	12-05-1999
			CZ 9802523 A	16-12-1998
			HU 9900621 A	30-08-1999
			WO 9730023 A	21-08-1997
			PL 328530 A	01-02-1999
WO 9909001	A	25-02-1999	AU 8562098 A	08-03-1999
			ZA 9807425 A	22-02-1999
WO 9318007	A	16-09-1993	AU 3648793 A	05-10-1993
EP 894787	A	03-02-1999	JP 9255637 A	30-09-1997
			AU 2176197 A	17-10-1997
			WO 9735835 A	02-10-1997
WO 9315041	A	05-08-1993	AT 156804 T	15-08-1997
			AU 3367993 A	01-09-1993
			CA 2107566 A	04-08-1993
			CN 1084846 A	06-04-1994
			DE 69313051 D	18-09-1997
			DE 69313051 T	08-01-1998
			DK 583485 T	06-10-1997
			EP 0583485 A	23-02-1994
			ES 2105286 T	16-10-1997
			GR 3025106 T	30-01-1998
			HK 1000768 A	24-04-1998
			HU 65351 A	02-05-1994
			IL 104567 A	18-03-1997
			JP 11092432 A	06-04-1999
			JP 2819435 B	30-10-1998
			JP 6506955 T	04-08-1994
			MX 9300552 A	01-09-1993
			RU 2125983 C	10-02-1999
			US 5387710 A	07-02-1995
			ZA 9300591 A	18-10-1993
WO 9218461	A	29-10-1992	AT 157080 T	15-09-1997
			DE 69221721 D	25-09-1997
			DK 579833 T	15-09-1997
			EP 0579833 A	26-01-1994
			JP 6506676 T	28-07-1994